

Middle Permian brachiopods from Setamai, the type locality of the Kanokura Formation, southern Kitakami Mountains, northeast Japan

Jun-ichi TAZAWA* and Yosuke IBARAKI**

Abstract

A Middle Permian (Kubergandian-Murgabian) brachiopod fauna is described from the type section of the lower Kanokura Formation in the Setamai area, southern Kitakami Mountains, northeast Japan. This fauna contains the following nine species: *Transennatia gratiosa*, *Tyloplecta* cf. *yangzeensis*, *Waagenoconcha* sp., *Linoproductus cora*, *Cancrinella* sp., *Leptodus nobilis*, *Derbyia grandis*, *Derbyia nipponica* and *Spiriferella keilhavii*. The Setamai fauna is characterized by the mixture of both the Boreal and Tethyan elements.

Key words: Boreal-Tethyan mixed fauna, brachiopods, Middle Permian, Setamai, southern Kitakami Mountains.

Introduction

The Permian brachiopod specimens described in this paper were collected by the authors and late Prof. M. Minato of Hokkaido University from nine localities in the Kanokurasawa and Kacchizawa valleys in the Setamai area, the type locality of the lower part of the Kanokura Formation, southern Kitakami Mountains, northeast Japan (Figs. 1, 2).

The Middle Permian Kanokura Formation was named by Onuki (1937) as the Kanokura Stage, but Onuki (1956) later changed the name to 'formation' with the outcrops along the Kanokurasawa valley as its type section. The stratigraphy of the Kanokura Formation in the Setamai area was described and discussed in detail by Minato et al. (1954, 1978, 1979), Onuki (1956, 1969), Murata (1964), Saito (1966, 1968) and Choi (1973, 1976). In palaeontology, many species of fusulinaceans (Choi, 1973), corals (Minato, 1955; Minato and Kato, 1965), brachiopods (Hayasaka, 1953; Hayasaka and Minato, 1956; Minato and Nakamura, 1956;

* Department of Geology, Faculty of Science, Niigata University, Niigata 950-2181, Japan

** Graduate School of Science and Technology, Niigata University, Niigata 950-2181, Japan
(Manuscript received 24 November, 2000; accepted 21 December, 2000)




Fig. 1. Index map showing the study area (Figs. 2, 3).

Nakamura et al., 1970) and ammonoids (Hayasaka, 1954) were described from the Kanokura Formation in this area. On brachiopods, the following six species have been hitherto described: *Chonetes (Plicochonetes) deplanata* (Waagen), *Linoproductus cora* (d'Orbigny) and *Derbyia magnifica* Licharew by Hayasaka and Minato (1956), *Permundaria asiatica* Nakamura, Kato and Choi by Nakamura et al. (1970), *Orthotetes rugosa* Fredericks by Minato and Nakamura (1956), and *Hamletella kitakamiensis* Hayasaka by Hayasaka (1953).

In this paper the brachiopod fauna of the lower Kanokura Formation in the Setamai area is described, and the age and palaeobiogeography of the Setamai fauna are discussed. The first author (J. T.) is responsible for the systematic descriptions and palaeobiogeography of the fauna, and the second author (Y. I.) is responsible for the field geology. All the specimens are housed in the Department of Geology, Faculty of Science, Niigata University.

Stratigraphy

In the Setamai area marine fossiliferous Permian rocks of continental shelf deposits are well developed, forming a synclinorium trending NNW-SSE and plunging towards the south




Fig. 2. Index map showing the fossil localities (KS1-9) in the Setamai area. Using the topographical map of "Setamai" scale 1:25,000 published by the Geographical Survey Institution of Japan.

(Figs. 3, 4). The Permian rocks in the Setamai area are divided into three formations, the Sakamotozawa, Kanokura and Toyoma Formations in ascending order (Fig. 5). The Lower Permian Sakamotozawa Formation is mainly composed of sandstone and conglomerate in the lower part and limestone in the upper part, about 600 m in total thickness. This formation covers the Lower Carboniferous Arisu and Odaira Formations by an angular unconformity (Minato, 1942). The Middle Permian Kanokura Formation consists of sandstone, argillaceous impure limestone and conglomerate in the lower part and limestone in the upper part, about 500 m in total thickness. This formation disconformably overlies the Sakamotozawa Formation (Minato et al., 1954). The Upper Permian Toyoma Formation is mainly made of black shale, with two conglomerate beds, over 500 m in total thickness. This formation conformably overlies




Fig. 3. Geological map of the Setamai area. 1: dyke rocks, 2: conglomerate of the Toyoma, Kanokura and Sakamotozawa Formations, 3: shale of the Toyoma Formation, 4: limestone of the upper Kanokura Formation, 5: argillaceous impure limestone of the lower Kanokura Formation, 6: sandstone of the lower Kanokura Formation, 7: limestone of the Sakamotozawa Formation, 8: shale of the Sakamotozawa Formation, 9: sandstone of the Sakamotozawa Formation, 10: undivided Lower Carboniferous, 11: conformity, 12: unconformity, 13: fault, 14: concealed fault.




Fig. 4. Cross sections of the Setamai area. Shading as for Fig. 3.

the Kanokura Formation.

The brachiopod fossils treated in this paper were collected from the sandstone, shale and argillaceous impure limestone of the upper part of the lower Kanokura Formation in nine localities (KS1-9). The fossil localities and horizons are shown in Figures 2, 5. A fusulinacean *Monodiexodina matsubaishi* (Fujimoto, 1956) often occurs together with the brachiopods from the sandstone and argillaceous impure limestone of the upper part of the lower Kanokura Formation.

The Setamai fauna

The brachiopods of Setamai described here, and the number of specimens are shown as follows:

<i>Transennatia gratiosa</i> (Waagen, 1884)	5
<i>Tyloplecta</i> cf. <i>yangzeensis</i> (Chao, 1927)	1
<i>Waagenoconcha</i> sp.	1
<i>Linoproductus cora</i> (d'Orbigny, 1842)	26
<i>Canocrinella</i> sp.	1
<i>Leptodus nobilis</i> (Waagen, 1883)	7
<i>Derbyia grandis</i> Waagen, 1884	2
<i>Derbyia nipponica</i> Nakamura, 1972	5
<i>Spiriferella keilhavii</i> (von Buch, 1846)	20

Among these species, *Waagenoconcha* sp., *Canocrinella* sp., *Derbyia grandis* and




Fig. 5. Columnar section of the Permian rocks in the Setamai area, showing the stratigraphic positions of the fossil localities (KS1-9). 1: shale, 2: sandstone, 3: conglomerate, 4: limestone of the upper Kanokura Formation, 5: argillaceous impure limestone of the lower Kanokura Formation, 6: limestone of the Sakamotozawa Formation, 7: conformity, 8: unconformity.

Spiriferella keilhavii are the Boreal or antitropical elements. On the other hand, *Transennatia gratiosa*, *Tyloplecta* cf. *yangzeensis* and *Leptodus nobilis* are the Tethyan elements. *Linoproductus cora* is a cosmopolitan species; and *Derbyia nipponica* is an endemic form. The Setamai fauna is certainly a Boreal-Tethyan mixed brachiopod fauna.

The age of the Setamai fauna is roughly assigned to the Middle Permian (Kungurian-lower Kazanian) by the occurrence of *Spiriferella keilhavii* (Waterhouse et al., 1978). The fossil-bearing sandstone and argillaceous impure limestone often contains a fusulinacean *Monodiexodina matsubaishi*, an index fossil of the lower Kanokura Formation (*Monodiexodina matsubaishi* Zone, Tazawa, 1976). The lower Kanokura Formation is correlated with the Middle Permian

(Ufimian-Kazanian) (Minato et al., 1978, 1979). From the above fossil evidence and stratigraphical position, we conclude that the upper part of the lower Kanokura Formation in the Setamai area can be correlated with the Middle Permian (Kubergandian-Murgabian).

Consequently, the Setamai fauna is a Middle Permian (Kubergandian-Murgabian) Boreal-Tethyan mixed brachiopod fauna in the southern Kitakami Mountains. This conclusion supports the Tazawa's opinion (Tazawa, 1991, 1998, 1999a, b) that the South Kitakami region was placed on the transitional zone between the Boreal and Tethyan Realms occupying the eastern sea of the Sino-Korean block, i.e., the Inner Mongolian-Japanese Transition Zone of Tazawa (1991) in the Middle Permian time.

Description of species

Order Productida Sarytcheva and Sokolskaya, 1959

Suborder Productidina Waagen, 1883

Superfamily Productoidea Gray, 1840

Family Productellidae Schuchert, 1929

Subfamily Marginiferinae Stehli, 1954

Tribe Paucispiniferini Muir-Wood and Cooper, 1960

Genus *Transennatia* Waterhouse, 1975

Transennatia gratiosa (Waagen, 1884)

Pl. 1, figs. 1-3.

Productus gratiosus Waagen, 1884, p. 691, pl. 72, figs. 3-7; Rothpletz, 1892, p. 76, pl. 10, figs. 15-15c; Diener, 1897, p. 23, pl. 3, figs. 3-7; Mansuy, 1913, p. 115, pl. 13, figs. 1a, b; Broili, 1916, p. 12, pl. 116, figs. 4, 5, 7-13; Colani, 1919, p. 10, pl. 1, figs. 2a-c; Chao, 1927, p. 44, pl. 4, figs. 6-10; Chi-Thuan, 1962, p. 491, pl. 2, figs. 5-7.

Productus (Dictyoclostus) gratiosus Waagen: Huang, 1933, p. 88, pl. 11, figs. 14a, b; Hayasaka, 1960, p. 49, pl. 1, fig. 8.

Marginifera gratiosa (Waagen): Reed, 1944, p. 98, pl. 19, figs. 6-7.

Dictyoclostus gratiosus (Waagen): Zhang and Ching (Jin), 1961, p. 411, pl. 4, figs. 12-18; Wang et al., 1964, p. 291, pl. 45, figs. 14-19.

Gratiosina gratiosa (Waagen): Grant, 1976, pl. 33, figs. 19-26; Licharew and Kotlyar, 1978, pl. 12, figs. 5, 6; pl. 20, figs. 1a, b; Minato et al., 1979, pl. 61, figs. 11-13.

Asioproductus gratiosus (Waagen): Yang et al., 1977, p. 350, pl. 140, figs. 5a-c; Feng and Jiang, 1978, p. 254, pl. 90, figs. 1-2; Tong, 1978, p. 228, pl. 80, figs. 7a, b; Lee et al., 1980, p. 373, pl. 164, figs. 14a-c; pl. 166, figs. 5-6.

Asioproductus bellus Chan (Zhan), 1979, p. 85, pl. 6, figs. 7-13; pl. 9, figs. 8-10; text-fig. 18.

Gratiosina sp. Minato et al., 1979, pl. 61, fig. 14; Tazawa, 1991, p. 215.

Transennatia graciosus (Waagen): Liu et al., 1982, p. 185, pl. 132, figs. 9a-d; Wang et al., 1982, p. 214, pl. 92, figs. 6-8; pl. 102, figs. 4-9; Ding and Qi, 1983, p. 280, pl. 95, figs. 14a, b.

Transennatia graciososa (Waagen): Yang, 1984, p. 219, pl. 33, figs. 7a-c; Jin, 1985, pl. 4, figs. 33, 34, 45, 46; Tazawa and Matsumoto, 1998, p. 6, pl. 1, figs. 4-8; Tazawa et al., 2000, p. 7, pl. 1, figs. 3-5.

Material.—Five specimens, from localities KS2, KS9: (1) external moulds of two ventral valves, NU-B302, 303; (2) internal moulds of two ventral valves, NU-B304, 305; (3) external mould of a dorsal valve, NU-B306.

Remarks.—These specimens are referred to *Transennatia graciososa* (Waagen, 1884), originally described from the Wargal and Chhidru Formations of the Salt Range, on the basis of their small shells, strongly convex ventral valve and reticulate ornament on the disks of both valves. The Setamai specimens are smaller (length 12 mm, width 13 mm in the largest specimen, NU-B305) than the Salt Range specimens, but similar to the shells of *T. graciososa* from the Middle Permian of the Hida Gaien Belt, central Japan (Tazawa, 1991; Tazawa and Matsumoto, 1998), the South Kitakami Belt, northeast Japan (Hayasaka, 1960; Minato et al., 1979; Tazawa et al., 2000), South Primorye, eastern Russia (Licharew and Kotlyar, 1978) and Heilongjiang and Jilin, northeast China (Lee et al., 1980).

Distribution.—Middle to Upper Permian: Salt Range, Himalayas (Gaqoi), Kumaon Himalayas, Cambodia, Vietnam, Timor, south China (Guangxi, Guangdong, Guizhou, Hunan, Sichuan and Hubei), northwest China (Shaanxi), east China (Jiangxi, Zhejiang and Anhui), northeast China (Jilin and Heilongjiang), eastern Russia (South Primorye) and Japan (Hida Gaien and South Kitakami Belts).

Family Productidae Gray, 1840

Subfamily Leioproductinae Muir-Wood and Cooper, 1960

Tribe Tyloplectini Termier and Termier, 1970

Genus *Tyloplecta* Muir-Wood and Cooper, 1960

Tyloplecta cf. *yangtzeensis* (Chao, 1927)

Pl. 1, fig. 5.

Compare.—

Productus yangtzeensis Chao, 1927, p. 50, pl. 5, figs. 1-3; pl. 8, fig. 9.

Productus (Dictyoclostus) yangtzeensis Chao: Huang, 1932, p. 26, pl. 1, figs. 1a-c.

Tyloplecta yangtzeensis (Chao): Schr ter, 1963, p. 124, pl. 6, figs. 1-7; Nakamura et al., 1981, p. 44, pl. 1, figs. 1-3; pl. 3, figs. 1, 2.

Material.—One specimen, from locality KS8, external mould of a dorsal valve and associated internal mould of a part of the dorsal valve, NU-B335.

Remarks.—This specimen is safely assigned to the genus *Tylopecta* by its large (length 48 mm, width about 52 mm), subquadrate dorsal valve with large ears, reticulate ornament on the posterior portion and a strong trilobate cardinal process. The Setamai specimen most resembles *T. yangtzeensis* (Chao, 1927), originally described from the Upper Permian of south and east China, in size, shape and external ornament of the dorsal valve. But the poor preservation of the present material makes accurate comparison difficult.

Tylopecta is a typical Tethyan-type genus and occurs mainly from the Lower to Middle Permian of south China.

Superfamily Echinoconchoidea Stehli, 1954

Family Echinoconchidae Stehli, 1954

Subfamily Juresaniinae Muir-Wood and Cooper, 1960

Tribe Waagenoconchini Muir-Wood and Cooper, 1960

Genus *Waagenoconcha* Chao, 1927

Waagenoconcha sp.

Pl. 1, fig. 4.

Material.—One specimen, from locality KS9, external mould of a dorsal valve and associated internal mould of a part of the dorsal valve, NU-B336.

Remarks.—The single incomplete dorsal valve specimen from Setamai is transverse subquadrate in outline, slightly concave, with low median fold, and strongly geniculated, length about 29 mm, width about 42 mm. The external surface of the dorsal valve is ornamented by numerous quincuncially arranged spine bases and a few weak concentric rugae. Only a strong trilobed cardinal process is observed as the dorsal valve internal structure. This specimen is safely assigned to the genus *Waagenoconcha* by its size, shape and external ornament of the dorsal valve, although the specific identification is difficult.

Waagenoconcha is a common antitropical genus in the Permian time (Tazawa, 1974, 1991; Shi et al., 1995).

Superfamily Linoproductoidea Stehli, 1954

Family Linoproductidae Stehli, 1954

Subfamily Linoproductinae Stehli, 1954

Genus *Linoproductus* Chao, 1927

Linoproductus cora (d'Orbigny, 1842)

Pl. 1, figs. 11-13; Pl. 2, figs. 1-8.

Productus cora, d'Orbigny, 1842, p. 55, pl. 5, figs. 8-10; Waagen, 1884, p. 677, pl. 66, fig. 3; pl. 67, figs. 1, 2; Schellwien, 1900, p. 41, pl. 7, figs. 15-17; Tschernyschew, 1902, p. 279, 621, pl. 33, figs. 2, 3; pl. 35, figs. 1a-c; pl. 54, figs. 1-5; text-figs. 69-71; Kozlowsky, 1914, p. 48, pl. 4, figs. 19a, b; pl. 5, fig. 5; pl. 6, figs. 1-10; Hayasaka, 1925, p. 94, pl. 5, figs. 7-9; Licharew, 1939, p. 88, pl. 19, figs. 5-7.

Productus (Productus) cora d'Orbigny: Licharew, 1937, p. 29, 101, pl. 6, figs. 1, 2.

Linoproductus cora (d'Orbigny): Chao, 1927, pars, p. 132, pl. 14, figs. 1a, b only; Ivanov, 1935, p. 105, pl. 5, fig. 4; Grabau, 1936, p. 123, pl. 12, figs. 7-9; Sarytcheva and Sokolskaya, 1952, p. 115, pl. 20, fig. 147; Chronic, 1953, p. 88, pl. 7, figs. 7-9; Hayasaka and Minato, 1956, p. 145, pl. 23, figs. 9, 10; Muir-Wood and Cooper, 1960, pl. 111, figs. 3-6; Zavodowsky et al., 1970, p. 94, pl. 5, fig. 3; pl. 36, fig. 2; pl. 62, figs. 10, 11; Samtleben, 1971, p. 80, pl. 7, figs. 2-5; text-fig. 13; Prokofiev, 1975, p. 32, pl. 4, figs. 1-2; Lee and Gu, 1976, p. 258, pl. 167, figs. 11, 12; Tazawa, 1976, pl. 2, fig. 11; Minato et al., 1979, pl. 62, figs. 1, 2; Lee et al., 1980, p. 376, pl. 152, figs. 6-8; pl. 169, fig. 10; Wang et al., 1982, p. 221, pl. 90, figs. 2, 4; Wang, 1984, p. 192, pl. 78, figs. 4a, b; Kalashnikov, 1986, pl. 119, fig. 1; Zhu, 1990, p. 77, pl. 15, figs. 21-24; Wang and Yang, 1998, p. 99, pl. 15, figs. 8-13.

Linoproductus hayasakai Tazawa, 1979, p. 26, pl. 4, figs. 5-11.

Linoproductus sp. Minato et al., 1979, pl. 62, figs. 3, 4.

Material.—Twenty-six specimens, from localities KS1, KS3, KS4: (1) external and internal moulds of four ventral valves, NU-B337-340; (2) internal moulds of nineteen ventral valves, NU-B341-359; (3) external moulds of three dorsal valves, NU-B360-362.

Description.—Shell large for genus, transverse subcircular in outline, widest at hinge; length about 56 mm, width about 80 mm in the largest specimen (NU-B341); length about 36 mm, width about 46 mm in the best preserved specimen (NU-B351). Ventral valve strongly convex in lateral profile, most convex at just posterior to midvalve and slightly convex to nearly flat on anterior half of valve, not geniculated. Anterior profile of the valve unevenly convex, with flattened venter and steeply inclined lateral slopes. Sulcus absent. Umbo small. Ears large, flattened, with blunt, angular extremities. Dorsal valve slightly concave on visceral disk, no fold. Trail not preserved. External surface of both valves ornamented by numerous, fine costellae and several strong rugae; costellae rather regularly developed, numbering 16-18 in 10 mm at midvalve; rugae occurring on ears of ventral valve and on whole surface of dorsal valve.

Internal structures of ventral valve are obscure in the present material.

Remarks.—These specimens are referred to *Linoproductus cora* (d'Orbigny, 1842), originally described from the Lower Permian of Apillapampa, Bolivia, in the trans-

verse shell outline and no ventral sulcus. The Setamai specimens have no spine bases on the ventral valves.

Linoproductus hayasakai Tazawa (1979, p. 26, pl. 4, figs. 5-11) and *Linoproductus* sp. (Minato et al., 1979, pl. 62, figs. 3, 4) from the lower Kanokura Formation of the southern Kitakami Mountains, also lack spine bases on the ventral valves, and they seem to be conspecific with *L. cora*.

Distribution.—Middle Carboniferous to Middle Permian: Peru, Bolivia (Apillapampa), western Russia (Moscow Basin), Urals, northern Russia (Timan), northeastern Russia (Kolyma), Carnic Alps, Caucasus, Uzbekistan (Fergana), Salt Range, south China (Guizhou and Hubei), east China (Fujian), northwest China (Gansu and Xinjiang), north China (Inner Mongolia), northeast China (Heilongjiang) and Japan (South Kitakami Belt).

Subfamily Grandaurispininae Lazarev, 1986

Genus *Cancrinella* Fredericks, 1928

Cancrinella sp.

Pl. 1, figs. 6a, b.

Material.—One specimen, from locality KS9, external mould of a ventral valve, NU-B307.

Remarks.—The single specimen from the Kanokurasawa valley is strongly deformed, but is safely assigned to the genus *Cancrinella* by its small size (length 17 mm, width 12 mm) and numerous elongate spine bases on the ventral valve. This specimen most resembles *Cancrinella kozlowskiana* (Fredericks, 1925, p. 18, pl. 1, figs. 36-40; pl. 2, figs. 86, 87) from the Middle Permian of Cape Kalouzin, South Primorye in size and external ornament. However, accurate comparison is difficult for the poorly preserved specimen.

Suborder Lyttoniida Williams, Harper and Grant, 2000

Superfamily Lyttonioidea Waagen, 1883

Family Lyttoniidae Waagen, 1883

Subfamily Lyttoniinae Waagen, 1883

Genus *Leptodus* Kayser, 1883

Leptodus nobilis (Waagen, 1883)

Pl. 1, figs. 7-10.

Lyttonia nobilis Waagen, 1883, p. 398, pl. 29, figs. 1-3; pl. 30, figs. 1, 2, 5, 6, 8, 10, 11; Diener, 1897, p. 37, pl. 1, figs. 5-7; Noetling, 1904, p. 112, text-figs. 4-7; Noetling, 1905, p. 140, pl. 17, figs. 1, 2; pl. 18, figs. 1-11; text-fig. 2; Mansuy, 1913, p. 123, pl. 13, fig. 10; Mansuy, 1914, p. 32, pl. 6, figs. 7a-d; pl. 7, figs. 1a-e; Diener, 1915, p. 99, pl. 10, fig. 15;

- Albrecht, 1924, p. 289, figs. 1a, b; Grabau, 1931, pars, p. 285, pl. 28, figs. 4, 5 only; Huang, 1932, p. 89, pl. 7, figs. 9, 10; pl. 8, figs. 8, 9; pl. 9, figs. 1-8; text-figs. 8-11; Simic, 1933, p. 49, pl. 4, fig. 1.
- Lyttonia tenuis* Waagen, 1883, p. 401, pl. 30, figs. 3, 4, 7, 9.
- Lyttonia* sp. Yabe, 1900, p. 2, text-figs. 1, 2.
- Lyttonia richthofeni* (Kayser): Frech, 1911, pars, p. 135, pl. 20, figs. 2a, b only; Hayasaka, 1917, p. 43, pl. 18, figs. 1-8; Hayasaka, 1922a, p. 62, pl. 11, figs. 1-6; Hayasaka, 1922b, p. 103, pl. 4, figs. 12, 13; Licharew, 1932b, p. 56, 86, pl. 1, figs. 1-16; pl. 2, figs. 1, 2, 5, 7, 10, 12; pl. 3, figs. 2-7; pl. 4, figs. 1-17; pl. 5, figs. 1-4, 6; Mashiko, 1934, p. 182, text-fig.
- Lyttonia* cf. *tenuis* Waagen: Mansuy, 1912, p. 19, pl. 4, fig. 4; pl. 5, figs. 1a-e; Huang, 1936, p. 493, pl. 1, fig. 6.
- Oldhamina* (*Lyttonia*) *richthofeni* var. *nobilis* Waagen: Fredericks, 1916, p. 76, pl. 4, fig. 2; text-fig. 22.
- Lyttonia* (*Leptodus*) *richthofeni* Kayser: Hamlet, 1928, p. 31, pl. 6, figs. 1-4.
- Lyttonia* cf. *richthofeni* (Kayser): Huang, 1932, p. 87, pl. 8, figs. 4a, b.
- Lyttonia richthofeni* Kayser forma *nobilis* Waagen: Licharew, 1932a, p. 69, 96, pl. 2, figs. 13, 14; pl. 5, figs. 1-4, 6; text-fig. 3.
- Leptodus nobilis* (Waagen): Wanner and Sieverts, 1935, p. 249, pl. 9, figs. 27, 28; text-figs. 16-18; Ramovs, 1958, p. 497, pl. 2, fig. 3; pl. 10, fig. 3; Termier and Termier, 1960, p. 241, pl. 3, figs. 1-10; Chi-Thuan, 1961, p. 274, pl. 1, figs. 1a, b; Schr ter, 1963, pl. 3, figs. 5-8; Sarytcheva, 1964, p. 65, pl. 7, figs. 5-8; text-fig. 1; Ruzhentsev and Sarytcheva, 1965, pl. 39, figs. 6-8; Cooper and Grant, 1974, pl. 191, figs. 8, 9; Grant, 1976, pl. 43, figs. 18, 19; Lee and Gu, 1976, p. 267, pl. 162, figs. 1, 2; Tazawa, 1976, pl. 2, fig. 8; Yang et al., 1977, p. 371, pl. 147, fig. 5; Feng and Jiang, 1978, p. 269, pl. 100, fig. 2; Licharew and Kotlyar, 1978, pl. 14, figs. 13-15; Jin et al., 1979, p. 82, pl. 23, fig. 15; Minato et al., 1979, pl. 66, figs. 1, 4, 5; Zhan, 1979, p. 93, pl. 9, fig. 12; Lee et al., 1980, p. 389, pl. 172, figs. 15, 16; Wang et al., 1982, p. 229, pl. 95, fig. 20; Zhan and Wu, 1982, pl. 4, fig. 4; Zhang et al., 1983, p. 297, pl. 102, figs. 7, 8; Yang, 1984, p. 226, pl. 35, fig. 12; Gu and Zhu, 1985, pl. 1, figs. 31, 33, 34; Liao and Meng, 1986, p. 81, pl. 2, figs. 24, 25; Sremac, 1986, p. 30, pl. 10, figs. 1-2; Tazawa, 1987, text-fig. 1.11; Kotlyar, in Kotlyar and Zakharov, 1989, pl. 20, fig. 6; pl. 23, fig. 12; Liang, 1990, p. 225, pl. 40, figs. 1, 5; Fang and Fan, 1994, p. 83, pl. 23, figs. 1-3; pl. 30, fig. 5; Tazawa and Matsumoto, 1998, p. 7, pl. 2, figs. 7-12; Tazawa et al., 1998, p. 241, figs. 2.1, 2.2, 4; Kato et al., 1999, p. 47, figs. 4a, b.
- Lyttonia* cf. *nobilis* Waagen: Huang, 1936, p. 493, pl. 1, fig. 5.
- Leptodus* cf. *nobilis* (Waagen): Thomas, 1957, p. 177, pl. 20, figs. 1-6.
- Leptodus richthofeni* Kayser: Shimizu, 1961, pl. 18, figs. 14, 15; Schr ter, 1963, p. 106, pl. 3, fig. 4; Sarytcheva, 1964, p. 65, pl. 7, figs. 2-4; Yang et al., 1977, p. 372, pl. 147, fig. 10; Yang, 1984, p. 226, pl. 35, fig. 11; Duan and Li, 1985, p. 119, pl. 35, figs. 17-19.
- Leptodus ivanovi* Fredericks: Minato et al., 1979, pl. 66, fig. 3.

Leptodus sp. Minato et al., 1979, pl. 66, fig. 2; Tazawa, 1987, text-fig. 1.10; Yanagida, 1996, fig. 2.14.

Leptodus tenuis (Waagen): Wang et al., 1982, p. 229, pl. 86, fig. 14; pl. 88, fig. 6; pl. 100, fig. 7; Duan and Li, 1985, p. 119, pl. 35, figs. 14-16; Liang, 1990, p. 226, pl. 40, fig. 9; Zhu, 1990, p. 79, pl. 18, figs. 19-21; Fang and Fan, 1994, p. 83, pl. 23, figs. 4-5; pl. 30, fig. 6.
Gubleria sp. Zhu, 1990, p. 80, pl. 16, fig. 24.

Material. — Seven specimens, from localities KS1, KS7, internal moulds of seven ventral valves, NU-B308-314.

Remarks. — These specimens are referred to *Leptodus nobilis* (Waagen, 1883), originally described from the Wargal and Chhidru Formations of the Salt Range, by their flat ventral valve with thick lateral ridges and nearly straight lateral lobes. The Setamai specimens are smaller than the type specimens. The dimensions of the best preserved specimen (NU-B308) with 18 lateral lobes are: length 34 mm, width 38 mm.

Leptodus richthofeni Kayser, 1883, originally described and figured by Kayser (1883, p. 161, pl. 21, figs. 9-11) from the Upper Permian of Jiangxi (Loping), South China and refigured by Cooper and Grant (1974, pl. 191, figs. 11-13), is easily distinguished from the present species by its more convex ventral valve with sharp lateral ridges and thick and strongly arched lateral lobes.

Distribution. — Lower Permian (Sakmarian) to Upper Permian (Dorashamian): southern Europe (Hungary, Croatia and Serbia), Transcaucasia, Pamir, Kashmir, Salt Range, Kumaon Himalayas, Tibet (Xizang), southern Thailand, Cambodia, Laos, Vietnam, Timor, northern Australia (Northern Territory), south China (Yunnan, Sichuan, Guizhou, Guangxi, Guangdong, Hunan and Hubei), east China (Jiangxi, Fujian, Zhejiang and Anhui), northwest China (Shaanxi, Gansu and Qinghai), north China (Inner Mongolia), northeast China (Heilongjiang and Jilin), eastern Russia (South Primorye) and Japan (South Kitakami, Hida Gaien, Maizuru, Mino and Akiyoshi Belts).

Order Orthotetida Waagen, 1884

Suborder Orthotetidina Waagen, 1884

Superfamily Orthotetoidea Waagen, 1884

Family Derbyiidae Stehli, 1954

Genus *Derbyia* Waagen, 1884

Derbyia grandis Waagen, 1884

Pl. 2, figs. 9a, b; Pl. 3, figs. 3a-c.

Derbyia grandis Waagen, 1884, p. 597, pl. 51, figs. 1-1c; pl. 52, figs. 1, 3; pl. 53, figs. 3, 5; Tschernyschew, 1902, p. 207, 580, pl. 24, figs. 1, 2; pl. 26, figs. 5a, b; text-figs. 59, 60;

Broili, 1916, p. 7, pl. 115, fig. 9; Frebold, 1950, p. 41, pl. 1, figs. 5, 5a; Grunt and Dmitriev, 1973, p. 84, pl. 3, figs. 1-4; Manankov, 1973, pl. 8, figs. 4, 5; Kulikov, 1974, p. 82, pl. 1, figs. 6a, b, v; Lee and Gu, 1976, p. 236, pl. 161, fig. 12; Licharew and Kotlyar, 1978, pl. 12, fig. 1; Lee et al., 1980, p. 336, pl. 159, fig. 13; Kalashnikov, 1986, pl. 114, fig. 1; Nakamura et al., 1992, pl. 1, fig. 1; Kalashnikov, 1993, p. 23, pl. 6, figs. 1, 2; pl. 7, fig. 7; Fang and Fan, 1994, p. 73, pl. 27, figs. 5-8; pl. 28, figs. 1, 2.

Derbyia cf. *grandis* Waagen: Frech, 1911, p. 125, pl. 18, figs. 4a-d; Stepanov, 1937, p. 110, 174, pl. 1, fig. 5; Harker, 1960, p. 52, pl. 16, figs. 9, 10; Bamber and Waterhouse, 1971, pl. 18, fig. 1; Manankov, 1979, p. 57, pl. 4, fig. 1; text-fig. 25.

Derbyia aff. *grandis* Waagen: Gobbett, 1963, p. 54, pl. 2, figs. 1, 2.

Material.—Two specimens, from localities KS1, KS9, external and internal moulds of two ventral valves, NU-B363, 364.

Description.—Shell large for genus, transverse outline; length about 63 mm, width 115 mm+ in the better preserved specimen (NU-B363). Ventral valve slightly concave to nearly flat in both profiles, ornamented by numerous costellae and several strong, irregular rugae. Costellae thin, with broad interspaces, and increasing by intercalation; numbering 10-11 in 10 mm at midvalve.

Ventral valve interior with strong, high median septum and large flabellate muscle scar. Median septum extending about one-third valve length. Muscle scar deeply impressed and surrounded by raised ridges.

Remarks.—These specimens can be referred to *Derbyia grandis* Waagen, 1884, originally described from the Wargal and Chhidru Formations of the Salt Range, on account of their large, transverse shells, and the external ornament of the ventral valves consisting of strong, irregular rugae and numerous costellae with broad interspaces. The type specimens from Salt Range differ from the present material in having more strongly convex ventral valve. The Setamai specimens closely resemble the shells, described as *Derbyia* cf. *grandis*, from the Assistance Formation of Devon Island (Harker, 1960, p. 52, pl. 16, figs. 9, 10) and from the Kapp Starostin Formation of Spitsbergen (Gobbett, 1963, p. 54, pl. 2, figs. 1, 2), in their slightly concave to nearly flattened ventral valve.

Distribution.—Lower to Upper Permian, mostly Lower and Middle Permian: Arctic Canada (northern Yukon Territory and Devon Island), northeast Greenland, Spitsbergen, northern Russia (Timan, northern Urals and Kolyma-Omolon region), Salt Range, southeast Pamir, Timor, southwest China (Yunnan), east China (Jiangsu), north China (Inner Mongolia), northeast China (Jilin), eastern Russia (South Primorye) and Japan (South Kitakami Belt).

Derbyia nipponica Nakamura, 1972

Pl. 3, figs. 1, 2, 4.

Derbyia magnifica Licharew: Hayasaka and Minato, 1956, p. 141, pl. 23, figs. 1a-c;
Hayasaka, 1960, p. 45, pl. 2, figs. 5, 6.

Derbyia nipponica Nakamura, 1972, p. 399, pl. 7, figs. 1, 4-9; Minato et al., 1979, pl. 60, figs. 1-4.

Material.—Five specimens, from localities KS1, KS9: (1) external and internal moulds of three ventral valves, NU-B365-367; (2) internal mould of a ventral valve, NU-B368; (3) internal mould of a dorsal valve, NU-B369.

Description.—Shell medium to large for genus, transversely subelliptical in outline; greatest width near midvalve; length 41 mm, width 49 mm in the best preserved ventral valve specimen (NU-B365); length 51 mm, width about 73 mm in the largest dorsal valve specimen (NU-B369).

Ventral valve slightly convex to nearly flat, ornamented by numerous fine costellae and 3-5 irregular, strong rugae. Costellae regularly developed, increasing by intercalation, and having narrow interspaces; numbering 19-20 in 10 mm at midvalve. Dorsal valve slightly but more strongly convex than ventral valve. External ornament obscure in the present material.

Internally, ventral valve having thin, long median septum, extending near midvalve. Dorsal valve having high cardinal process and divergent, strong crural plates.

Remarks.—The Setamai specimens are referred to *Derbyia nipponica* Nakamura, 1972, originally described as *Derbyia magnifica* Licharew, 1932, by Hayasaka and Minato (1956, p. 141, pl. 23, figs. 1a-c) from the lower Kanokura Formation in the Imo and Setamai areas, southern Kitakami Mountains, and redescribed as *D. nipponica* by Nakamura (1972, p. 399, pl. 7, figs. 1, 4-9) on new materials from the Imo area, one of the type localities. *D. nipponica* is characterized by its weakly biconvex and transverse shell, short hinge, and regularly developed, fine costellae with narrow interspaces.

Derbyia magnifica Licharew, 1932, originally described as *Derbyia magna* Licharew (1932a, p. 18, 40, pl. 1, figs. 1, 2, 6, 7, 13; pl. 2, figs. 1a-c; pl. 3, figs. 1a, b) from the Lower Permian of northern Caucasus, differs from *D. nipponica* in having long hinge marking the greatest width, ventral fold and dorsal sulcus.

Derbyia regularis Waagen (1884, p. 594, pl. 53, figs. 1, 2, 4) from the Amb and Wargal Formations of the Salt Range is also medium to large, transverse *Derbyia*, but it differs from the present species in having more less dense costellae.

Distribution.—Middle Permian of Japan (South Kitakami Belt).

Order Spiriferida Waagen, 1883
Suborder Spiriferidina Waagen, 1883
Superfamily Spiriferoidea King, 1846
Family Spiriferellidae Waterhouse, 1968
Subfamily Spiriferellinae Waterhouse, 1968
Genus *Spiriferella* Tschernyschew, 1902

Spiriferella keilhavii (von Buch, 1846)

Pl. 4, figs. 1-10.

Spirifer keilhavii von Buch, 1846, p. 74, figs. 2a, b.*Spiriferina keilhavii* (von Buch): Wiman, 1914, pl. 2, figs. 25-30; pl. 3, fig. 1.*Spiriferina draschei* (Toula): Wiman, 1914, p. 38, pl. 3, fig. 2 only.

Spiriferella keilhavii (von Buch): Frebold, 1931, p. 28, pl. 5, figs. 7-9; Grabau, 1931, p. 164, pl. 20, figs. 9a-c; pl. 21, figs. 1-5; Frebold, 1937, p. 46, pl. 11, fig. 9; Stepanov, 1937, p. 143, 179, pl. 7, figs. 8-11; Dunbar, 1955, p. 139, pl. 25, figs. 1-9; pl. 26, figs. 1-11; pl. 27, figs. 1-14; Harker, 1960, p. 72, pl. 22, figs. 9-11; pl. 23, figs. 1, 2; Gobbett, 1963, p. 152, pl. 20, figs. 8-10; Yanagida, 1963, pl. 9, figs. 4-9; pl. 10, figs. 1-7; Nelson and Johnson, 1968, p. 736, pl. 96, figs. 7, 8, 12; text-figs. 3e, 8a, 9, 13b; Koizumi, 1979, pl. 1, figs. 10, 11 only; Minato et al., 1979, pl. 67, figs. 1-3; Lee et al., 1980, p. 418, pl. 178, fig. 8; Tazawa and Gunji, 1982, p. 70, pl. 4, figs. 4-7; Waterhouse and Waddington, 1982, p. 28, pl. 6, figs. 3-14; text-figs. 16e, g, h, i, 19; not pl. 4, fig. 15; Lee et al., 1985, p. 122, pl. 2, figs. 1a, b, 5, 8; Tazawa et al., 1999, p. 160, fig. 2.2-5; Tazawa, 1999a, p. 5, pl. 1, figs. 2-6.

Spiriferella parryana (Toula): Frebold, 1937, p. 45, pl. 11, fig. 6.*Spiriferella* aff. *keilhavii* (von Buch): Bamber and Waterhouse, 1971, pl. 20, fig. 7.

Material.—Twenty specimens, from locality KS9: (1) external and internal moulds of two ventral valves, NU-B315, 316; (2) external moulds of two ventral valves, NU-B317, 318; (3) external mould of a dorsal valve and internal mould of the conjoined valve, NU-B319; (4) external and internal moulds of a dorsal valve, NU-B320; (5) external moulds of four dorsal valves, NU-B321-324; (6) internal mould of a conjoined valve, NU-B325; (7) internal moulds of six ventral valves, NU-B326-331; (8) internal moulds of three dorsal valves, NU-B332-334.

Description.—Shell large for genus, slightly transverse subquadrate in outline, with greatest width at or near hinge; length more than 29 mm, width about 63 mm in the largest specimen (NU-B317); length 29 mm, width about 30 mm in the best preserved ventral valve specimen (NU-B318). Ventral valve gently convex in lateral profile, with maximum convexity at umbonal slope. Beak and interarea not well preserved. Sulcus wide and deep, with U-shaped cross-section and two minor costae on each side of inner slopes. External surface of ventral valve ornamented by 10-12 strong costae and numerous very fine concentric lirae; costae broad, rounded, and projecting fascicles of 2-3. Dorsal valve slightly convex in lateral profile. Fold wide and low, with a deep median groove commencing at hinge. External ornament of dorsal valve same as ventral valve.

Internally, ventral valve having a deeply impressed, large, heart-shaped muscle field and a pair of strong dental plates; adductors long and narrow; diductors wide and weakly striated longitudinally. Dorsal valve having a small, striated cardinal process. Other internal structures

Remarks.—These specimens are referred to *Spiriferella keilhavii* (von Buch, 1846), originally described from the Middle Permian of Mt. Misery, Bear Island, by their transverse shell, weakly fasciculated costae on both valves and broad, low fold with a deep median groove.

Spiriferella lita Fredericks, 1924, from the Chandalaz Formation of South Primorye (Fredericks, 1924, p. 36, pl. 1, figs. 16-27, text-fig. 2) differs from *S. keilhavii* in having strong and simple costae on the ventral valve.

Spiriferella loveni (Diener, 1903), redescribed and refigured by Waterhouse and Waddington (1982, p. 22, pl. 5, figs. 2-17; pl. 6, figs. 1, 2; text-figs. 16b, d, f, 17, 18) on the specimens from the Middle Permian of the Canadian Arctic region, is distinguished from *S. keilhavii* by its high, distinct dorsal fold with a broad, deep median groove.

Distribution.—Middle Permian (Bolorian-Midian): Arctic Canada (Yukon Territory, Devon Island and Ellesmere Island), Greenland, Svalbard (Spitsbergen and Bear Island), north China (Inner Mongolia), northeast China (Heilongjiang) and Japan (South Kitakami and Hida Gaien Belts).

Acknowledgements

We thank Dr. I. Niikawa of the Department of Geology, Niigata University and Dr. G. R. Shi of the School of Ecology and Environment, Deakin University for critical reading of the manuscript; Dr. K. Nakamura of Sapporo for providing a part of the specimens from the Minato collection stored in the Department of Earth and Planetary Sciences, Hokkaido University.

References

- Albrecht, J., 1924, Paläontologische und stratigraphische Ergebnisse der Forschungsreise nach Westserbien 1918. *A. Wiss. Wien, Math.-Naturwiss.*, **99**, 289-307.
- Bamber, E.W. and Waterhouse, J.B., 1971, Carboniferous and Permian stratigraphy and paleontology, northern Yukon Territory, Canada. *Bull. Canad. Petrol. Geol.*, **19**, 29-250.
- Broili, F., 1916, Die permischen Brachiopoden von Timor. *Paläont. Timor*, **7**, 1-104.
- Buch, L. von, 1846, Ueber *Spirifer keilhavii*. *Abh. k. Akad. Wiss. Berlin*, 1846, 65-80.
- Chao, Y.T., 1927, Productidae of China, pt. 1. Producti. *Palaeontologia Sinica, Ser. B*, **5**, 1-244.
- Chi-Thuan, T., 1961, Les brachiopodes permien du Phnom-Tup (Sisophon-Cambodge). *Ann. Fac. Sci. Saigon*, 1961, 267-308.
- Chi-Thuan, T., 1962, Les brachiopodes de Cam-Lo (Province de Quang-Tri). *Ann. Fac. Sci. Saigon*, 1962, 485-498.
- Choi, D.R., 1973, Permian fusulinids from the Setamai-Yahagi district, southern Kitakami Mountains, N. E. Japan. *Jour. Fac. Sci., Hokkaido Univ., Ser. 4*, **16**, 1-132.
- Choi, D.R., 1976, Distribution of the Upper Permian fusulinids with relation to limestone lithofacies in the southern Kitakami Mountains, N. E. Japan. *Jour. Geol. Soc. Japan*, **82**,

- lithofacies in the southern Kitakami Mountains, N. E. Japan. *Jour. Geol. Soc. Japan*, **82**, 113-125.
- Chronic, J., 1953, Systematic paleontology: Brachiopoda. In Newell, N.D., Chronic, J. and Roberts, T.G., Upper Paleozoic of Peru. *Geol. Soc. Amer. Mem.*, **58**, 49-108.
- Colani, M.M., 1919, Sur quelques fossiles Ouralo-Permians de Hongay. *Bull. Serv. Géol. l'Indochine*, **6**, 1-27.
- Cooper, G.A. and Grant, R.E., 1974, Permian brachiopods of west Texas, 2. *Smithson. Contr. Paleobiol.*, no. 15, 233-793.
- Diener, C., 1897, Himalayan fossils, vol. 1, pt. 3. The Permocarboniferous fauna of Chitichun, No. 1. *Palaeontologia Indica, Ser. 15*, **1**, pt. 3, 1-105.
- Diener, C., 1903, Himalayan fossils, vol. 1, pt. 5. Permian fossils of the central Himalayas. *Palaeontologia Indica, Ser. 15*, **1**, pt. 5, 1-204.
- Diener, C., 1915, The Anthracolithic faunae of Kashmir, Kanaur and Spiti. *Palaeontologia Indica, N. S.*, **5**, 1-135.
- Ding, P. and Qi, W., 1983, Phulum Brachiopoda. In Xian Institute of Geology and Mineral Resources, ed., *Palaeontological atlas of northeast China: Shaanxi, Gansu and Ningxia volume, pt. 2. Upper Palaeozoic*, Geol. Pub. House, Beijing, 244-425. (in Chinese)
- Duan, C. and Li, W., 1985, Descriptions of fossils; Phylum Brachiopoda. In Ding, Y., Xia, G., Duan, C. and Li, W., Study on the Early Permian stratigraphy and fauna in Zhesi district, Nei Mongol Zizhiqu (Inner Mongolia). *Bull. Tianjin Inst. Geol. Min. Res., Chinese Acad. Geol. Sci.*, no. 10, 99-145, 199-214. (in Chinese)
- Dunbar, C.O., 1955, Permian brachiopod faunas of central East Greenland. *Meddel. Grønland*, **110**, 1-169.
- Fang, R. and Fan, J., 1994, *Middle to Upper Carboniferous-Early Permian Gondwana facies and palaeontology in western Yunnan*. Yunnan Sci. Tech. Press, Kunming, 121 p. (in Chinese)
- Feng, R. and Jiang, Z., 1978, Phylum Brachiopoda. In Geological and Palaeontological Team of Guizhou, ed., *Palaeontological atlas of southwest China; Guizhou, pt. 2. Carboniferous to Quaternary*, Geol. Pub. House, Beijing, 231-305. (in Chinese)
- Frebold, H., 1931, Das marine Oberkarbon Ostgrönlands. Leitende Fauna, Altersstellung, Palaeogeographie. *Meddel. Grønland*, **84**, 1-88.
- Frebold, H., 1937, Das Festungsprofil auf Spitzbergen, 4. Die Brachiopoden- und Lamellibranchiatenfauna und die Stratigraphie des Oberkarbons und Unterperms. *Skrift. Svalbard og Ishavet*, no. 69, 1-94.
- Frebold, H., 1950, Stratigraphie und Brachiopodenfauna des marinen Jungpalaeozoikums von Holms und Amdrups Land (Nordostgrönland). *Mus. Minéral. Géol. l'Univ. Copenhague, Com. Géol.*, no. 37, 1-97.
- Frech, F., 1911, Die Dyas. In Richthofen, F. von, ed., *China, Fünfter Band*, Dietrich Reimer, Berlin, 103-137.
- Fredericks, G., 1916, The palaeontological notes, 2. On some Upper Palaeozoic Brachiopoda of Eurasia. *Mém. Com. Géol., N. S.*, **156**, 1-87. (in Russian)
- Fredericks, G., 1924, Ussuriyskiy verkhniy paleozoy, 1. Brachiopoda. *Mater. Geol. Polezn. Iskopaem. Dalnego Vostoka*, no. 28, 1-52. (in Russian)
- Fredericks, G., 1925, Ussuriyskiy verkhniy paleozoy, 2. Permskie brachiopody s mysa Kaluzina. *Mater. Geol. Polezn. Iskopaem. Dalnego Vostoka*, no. 40, 1-28. (in Russian)
- Fredericks, G., 1928, Materialy dlya klassifiatsii roda *Productus* Sow. *Izv. Geol. Kom., Leningrad*, **46**, 773-792. (in Russian)
- Fujimoto, H., 1956, A new species of *Parafusulina* from the Kitakami Massif, Japan. *Trans. Proc. Palaeont. Soc. Japan, N. S.*, no. 21, 157-160.
- Gobbett, D.J., 1963, Carboniferous and Permian brachiopods of Svalbard. *Norsk Polarinst. Skrift.*, no. 127, 1-201.

- Grabau, A.W., 1931, The Permian of Mongolia. In Reeds, C.A., ed., *Natural history of Central Asia*, vol. 4. Amer. Mus. Nat. Hist., New York, 665 p.
- Grabau, A.W., 1936, Early Permian fossils of China, pt. 2. Fauna of the Maping Limestone of Kwangsi and Kweichow. *Palaeontologia Sinica, Ser. B*, **8**, 1-441.
- Grant, R.E., 1976, Permian brachiopods from southern Thailand. *Jour. Paleontology*, **50**, 1-269.
- Gray, J.E., 1840, *Synopsis of the contents of the British Museum*, 42nd edit. London, 370 p.
- Grunt, T.A. and Dmitriev, V.Yu., 1973, Permskie brachiopody Pamira. *Tr. Paleont. Inst.*, **136**, 1-209. (in Russian)
- Gu, F. and Zhu, R., 1985, Lower Permian brachiopods from Lin-Dong, Nei Mongol. *Bull. Shenyang Inst. Geol. Min. Res., Chinese Acad. Geol. Sci.*, no. 12, 74-97. (in Chinese)
- Hamlet, B., 1928, Permische Brachiopoden, Lamellibranchiaten und Gastropoden von Timor. *Jaar. Mijnw. Ned.-Indie*, **56**, 1-115.
- Harker, P., 1960, Corals, brachiopods and molluscs of Grinnell Peninsula. In Harker, P. and Thorsteinsson, R., Permian rocks and faunas of Grinnell Peninsula, Arctic Archipelago. *Geol. Surv. Canada, Mem.*, **309**, 39-79.
- Hayasaka, I., 1917, On the brachiopod genus *Lyttonia* with several Japanese and Chinese examples. *Jour. Geol. Soc. Tokyo*, **24**, 43-53.
- Hayasaka, I., 1922a, Some Permian brachiopods from the Kitakami Mountains. *Japan. Jour. Geol. Geogr.*, **1**, 51-70.
- Hayasaka, I., 1922b, Paleozoic Brachiopoda from Japan, Korea and China. *Sci. Rep., Tohoku Imp. Univ., 2nd Ser.*, **6**, 1-116.
- Hayasaka, I., 1925, On some brachiopods from the *Lyttonia* horizon of the Kitakami Mountains. *Japan. Jour. Geol. Geogr.*, **4**, 89-103.
- Hayasaka, I., 1953, *Hamletella*, a new Permian genus of Brachiopoda and a new species from the Kitakami Mountains, Japan. *Trans. Proc. Palaeont. Soc. Japan, N. S.*, no. 12, 89-95.
- Hayasaka, I., 1954, Younger Palaeozoic cephalopods from the Kitakami Mountains. *Jour. Fac. Sci., Hokkaido Univ., Ser. 4*, **8**, 361-374.
- Hayasaka, I., 1960, On the occurrence of *Neospirifer fasciger* (Keyserling) in Japan, and a note on some associate Permian brachiopods from around Kesenuma City, northeast Japan. *Coll. Essays Comm. 10th Anniv. Shimane Univ.*, 34-57.
- Hayasaka, I. And Minato, M., 1956, Some brachiopods from the Lower Kanokura Series of the Kitakami Mountains, Japan. *Trans. Proc. Palaeont. Soc. Japan, N. S.*, no. 21, 141-147.
- Huang, T.K., 1932, Late Permian Brachiopoda of southwestern China. *Palaeontologia Sinica, Ser. B*, **9**, fasc. 1, 1-138.
- Huang, T.K., 1933, Late Permian Brachiopoda of southwestern China, pt. 2. *Palaeontologia Sinica, Ser. B*, **9**, fasc. 2, 1-172.
- Huang, T.K., 1936, On the occurrence of *Lyttonia* in the Wolfcamp Series of the Glass Mountains of Texas with notes on lyttonids from southwestern China. *Bull. Geol. Soc. China*, **15**, 489-493.
- Ivanov, A.P., 1935, Fauna brachiopod srednego i verkhnego karbona Podmoskovnogo basseyna, chasti 1. Productidae Gray, vypsik 1. Productinae Schuchert emend. Paeckelmann. *Tr. Mosk. Geol. Trests*, fasc. 8, 1-163. (in Russian)
- Jin, Y., 1985, Permian Brachiopoda and Paleogeography of the Qinghai-Xizang (Tibet) Plateau. *Palaeont. Cathayana*, no. 2, 19-71.
- Jin, Y., Ye, S., Xu, H. and Sun, D., 1979, Phylum Brachiopoda. In Nanjing Institute of Geology and Palaeontology, Academia Sinica and Qinghai Institute of Geology, eds., *Palaeontological atlas of northwest China; Qinghai volume, pt. 1*, Geol. Pub. House, Beijing, 60-217. (in Chinese)
- Kalashnikov, N.V., 1986, Brachiopody. In Gorskii, V.P. and Kalmykova, M.A., eds., Atlas kharakternykh kompleksov permskoi fauny i flory Urala: Russkoi platformy, *Tr. VSEGEI*, **331**, 29-30, 89-94. (in Russian)

- Kalashnikov, N.V., 1993, *Brakhiopody permi Evropeyskogo Severa Rossii*. Nauka, St. Peterburg, 151 p. (in Russian)
- Kato, M., Takeuchi, K., Hendarsyah, A. and Sundari, D., 1999, On the occurrence of the Permian brachiopod genus *Leptodus* in Timor. *Geol. Res. Develop. Contr., Bandung, Paleontology Series*, no. 9, 43-51.
- Kayser, E., 1883, Obercarbonische Fauna von Lo-Ping. In Richthofen, F. von, ed., *China, Vierter Band*, Dietrich Reimer, Berlin, 160-208.
- King, W., 1846, Remarks on certain genera belonging to the class Palliobranchiata. *Ann. Mag. Nat. Hist.*, **18**, 26-42.
- Koizumi, H., 1979, Permian brachiopods from Takakura-yama Formation and its geological age. N. E. Japan. *Bull. Taira Chigaku Dokokai, Spec. Vol.*, 1-3. (in Japanese)
- Kotlyar, G.V. and Zakharov, Yu. D., 1989, *Pozdnepermiskiy etap evolyutsii organicheskogo mira; Midiyskiy yarus SSSR*. Nauka, Leningrad, 182 p. (in Russian)
- Kozłowski, R., 1914, Les brachiopodes du Carbonifère supérieur de Bolivie. *Ann. Paléontologie*, **9**, 1-100.
- Kulikov, M.V., 1974, Brakhiopody kungurskogo yarusa Urala. *Tr. Inst. Geol. Geokhim., Ural. Nauch. Tsentr, Akad. Nauk SSSR*, no. 109, 77-123. (in Russian)
- Lazarev, S.S., 1986, Osnovnye napravleniya evolyutsii i sistema brakhiopod podotryada Productidina. *Avtoref. dis. d-ra biol., Nauk. Moskva*, 41 p. (in Russian)
- Lee, L. and Gu, F., 1976, Carboniferous and Permian Brachiopoda. In Geological Bureau of Nei Mongol and Geological Institute of northeast China, eds., *Palaeontological atlas of northeast China; Nei Mongol, pt. 1. Palaeozoic volume*, Geol. Pub. House, Beijing, 228-306. (in Chinese)
- Lee, L., Gu, F. and Li, W., 1985, *Spiriferella* and *Spiriferelloides* (Brachiopoda) from the Lower Permian of Xi Ujimqin Qi region, Inner Mongolia. *Prof. Pap. Stratigraphy and Paleontology*, no. 12, 121-139. (in Chinese)
- Lee, L., Gu, F. and Su, Y., 1980, Carboniferous and Permian Brachiopoda. In Shenyang Institute of Geology and Mineral Resources, ed., *Palaeontological atlas of northeast China, pt. 1. Palaeozoic volume*, Geol. Pub. House, Beijing, 327-428. (in Chinese)
- Liang, W., 1990, Lengwu Formation of Permian and its brachiopod fauna in Zhejiang Province. *Geol. Mem. Minist. Geol. Min. Res., P. R. China, Ser. 2*, no. 10, 1-522. (in Chinese)
- Liao, Z. and Meng, F., 1986, Late Chanxianian brachiopods from Huatang of Xian County, southern Hunan. *Mem. Nanjing Inst. Geol. Palaeont., Acad. Sinica*, no. 22, 71-94. (in Chinese)
- Licharew, B.K., 1932a, Fauna permskikh otlozheniy Severnogo Kavkaza, 1. Brachiopoda, Podsem. Orthotetinae Waagen. *Tr. VGRO*, fasc. 215, 3-54. (in Russian)
- Licharew, B.K., 1932b, Fauna permskikh otlozheniy Severnogo Kavkaza, 2. Brachiopoda, Sem. Lytoniidae Waagen. *Tr. VGRO*, fasc. 215, 55-84. (in Russian)
- Licharew, B.K., 1937, *Monografiya po paleontologii SSSR, tom 39. Brakhiopody permskikh otlozheniy SSSR, bypusk 1. Permskie Brachiopoda sebernogo Kavkaza, Semeystava: Chonetidae Hall et Clarke i Productidae Gray*. ONTI NKTP SSSR, Leningrad, 152 p. (in Russian)
- Licharew, B.K., 1939, Klass Brachiopoda. In Gorsky, I., ed., *Atlas rukovodyaschikh form iskopaemykh faun SSSR, tom 5. Spodniy i verkhniy otdely kamennougolnoy sistemy*, GONTI, Leningrad, 79-113. (in Russian)
- Licharew, B.K. and Kotlyar, G.V., 1978, Permskie brakhiopody Yuzhnogo Primorya. In Popeko, L.I., ed., *Verkhniy Paleozoy Severo-Vostochnoy Azii, Dalnovpstoch. Nauch. Chentr, Akad. Nauk SSSR, Vladivostok*, 63-75. (in Russian)
- Liu, Z., Tan, Z. and Ding, Y., 1982, Phylum Brachiopoda. In Geological Bureau of Hunan, ed., *The Palaeontological atlas of Hunan*, Geol. Pub. House, Beijing, 172-216. (in Chinese)

- Manankov, I.N., 1973, K pevizii rodov *Orthotetes* i *Derbyia* (Brachiopoda). *Paleontologicheskii Zhurnal*, 1973, no. 2, 52-60. (in Russian)
- Manankov, I.N., 1979, Strofomenidy pozdnego paleozoya Sibiri i Arktiki. *Tr. Paleont. Inst.*, **171**, 1-101. (in Russian)
- Mansuy, H., 1912, Mission du Laos, 1. Géologie des environs de Luang-Prabang. *Mém. Serv. Géol. l'Indochine*, **1**, 1-32.
- Mansuy, H., 1913, Faunes des calcaires a productus de l' Indochine, Première serie. *Mém. Serv. Géol. l'Indochine*, **2**, 1-132.
- Mansuy, H., 1914, Faunes des calcaires a productus de l'Indochine, Deuxième série. *Mém. Serv. Géol. l'Indochine*, **3**, 1-59.
- Mashiko, K., 1934, Discovery of *Lyttonia* in a limestone exposed at Takauti, Nakayakunomura, Amata-gun, Kyoto Prefecture. *Japan. Jour. Geol. Geogr.*, **11**, 181-183.
- Minato, M., 1942, Unconformity of the pre-Sakamotozawa Stage (pre-Sakmarian) in the Kitakami Mountainland, northeast Japan. *Jour. Geol. Soc. Japan*, **49**, 47-72. (in Japanese)
- Minato, M., 1955, Japanese Carboniferous and Permian corals. *Jour. Fac. Sci., Hokkaido Univ., Ser. 4*, **9**, 1-202.
- Minato, M., Hashimoto, S., Suyama, K., Takeda, Y., Suzuki, Y., Kimura, S., Yamada, K., Kakimi, T., Ichikawa, T. and Suetomi, H., 1954, Zur Biostratigraphie der permischen Formation des Setamai-Geländes im Süd-Kitakami Gebirge. *Jour. Geol. Soc. Japan*, **60**, 378-387. (in Japanese)
- Minato, M., Hunahashi, M., Watanabe, J. and Kato, M., 1979, *Variscan geohistory of northern Japan: The Abean Orogeny*. Tokai Univ. Press, Tokyo, 427 p.
- Minato, M. and Kato, M., 1965, Waagenophyllidae. *Jour. Fac. Sci., Hokkaido Univ., Ser. 4*, **12**, 1-241.
- Minato, M., Kato, M., Nakamura, K., Hasegawa, Y., Choi, D.R. and Tazawa, J., 1978, Biostratigraphy and correlation of the Permian of Japan. *Jour. Fac. Sci., Hokkaido Univ., Ser. 4*, **18**, 11-47.
- Minato, M. and Nakamura, K., 1956, On four brachiopod species of the subfamily Orthotetinae from the Kanokura Series of the Kitakami Mountains, Japan. *Trans Proc. Palaeont. Soc. Japan, N. S.*, no. 21, 149-155.
- Muir-Wood, H.M. and Cooper, G.A., 1960, Morphology, classification and life habits of the Productoidea (Brachiopoda). *Geol. Soc. Amer., Mem.*, **81**, 1-135.
- Murata, M., 1964, Geological age of the Kanokura Formation in the southern part of the Kitakami Massif, northeast Japan. *Saito Ho-on Kai Mus. Res. Bull.*, no. 33, 17-29.
- Nakamura, K., 1972, Permian Davidsoniacea from the southern Kitakami Mountains, Japan. *Jour. Fac. Sci., Hokkaido Univ., Ser. 4*, **15**, 361-426.
- Nakamura, K., Kato, M. and Choi, D.R., 1970, On *Permundaria*, a new genus of the brachiopod family Linoproductidae. *Jour. Fac. Sci., Hokkaido Univ., Ser. 4*, **14**, 293-299.
- Nakamura, K., Shimizu, D. and Golshani, F., 1981, On the occurrence of *Tyloplecta* in the Permian of the Abadeh region, central Iran. *Geol. Surv. Iran, Rep.*, no. 49, 39-53.
- Nakamura, K., Tazawa, J. and Kumon, F., 1992, Permian brachiopods of the Kapp Starostin Formation, west Spitsbergen. In Nakamura, K., ed., *Investigations on the Upper Carboniferous-Upper Permian succession of west Spitsbergen 1989-1991*, Hokkaido Univ., Sapporo, 77-95.
- Nelson, S.J. and Johnson, C.E., 1968, Permo-Pennsylvanian brachythyrid and horridonid brachiopods from the Yukon Territory, Canada. *Jour. Paleontology*, **42**, 715-746.
- Noetling, F., 1904, Ueber den Bau und die Organisation der Lytoniidae Waagen. *Verhandl. Deut. Zool. Gesell.*, 1904, 103-122.
- Noetling, F., 1905, Untersuchungen über die Familie Lytoniidae Waag. emend. Noetling. *Palaeontographica*, **51**, 129-154.
- Onuki, Y., 1937, On the Palaeozoic formation near Sakamotozawa, Kitakami Mountainland.

- Jour. Geol. Soc. Japan*, **44**, 168-186. (in Japanese)
- Onuki, Y., 1956, Geology of the Kitakami Massif. In Iwate Prefecture, ed., *Explanatory text of the geology of Iwate Prefecture*. Sasaki Shuppan, Sendai, 1-189. (in Japanese)
- Onuki, Y., 1969, Geology of the Kitakami Massif, northeast Japan. *Contr. Inst. Geol. Paleont. Tohoku Univ.*, no. 69, 1-239. (in Japanese)
- Orbigny, A. d', 1842, *Voyages dans l'Amerique Méridionale de 1826-1833, tome 3, pt. 4. Paléont.*, 188 p.
- Prokofiev, V.A., 1975, Brakhiopody verkhnego karbona Samarskoy luki. *Tr. Bsesoyuz. Nauch.-Issled. Geol. Razved Neft. Inst.*, no. 162, 1-144. (in Russian)
- Ramovs, A., 1958, Razvoj zgornjega perma v Loskih in Polhograjskih hribih. *Razprave, Dissertationes*, **4**, 451-622.
- Reed, F.R.C., 1944, Brachiopoda and Mollusca from the Productus Limestone of the Salt Range. *Palaeontologia Indica, N. S.*, **23**, 1-596.
- Rothpletz, A., 1892, Die Perm-, Trias- und Jura-Formation auf Timor und Rotti im indischen Archipel. *Palaeontographica*, **39**, 57-106.
- Ruzhentsev, V.E. and Sarytcheva, T.G., 1965, Razvitie i smena morskikh organizmov na rubezhe Paleozoya i Mesozoya. *Tr. Paleont. Inst.*, **108**, 1-431. (in Russian)
- Saito, Y., 1966, Geology of the Setamai district, southern Kitakami Massif, northeast Japan. *Contr. Inst. Geol. Paleont. Tohoku Univ.*, no. 62, 55-67. (in Japanese)
- Saito, Y., 1968, Geology of the Younger Paleozoic System of the southern Kitakami Massif, Iwate Prefecture, Japan. *Sci. Rep. Tohoku Univ., 2nd Ser.*, **40**, 79-139.
- Samtleben, C., 1971, Zur Kenntnis der Produktiden und Spiriferiden des bolivianischen Unterperms. *Beih. Geol. Jb.*, **111**, 1-163.
- Sarytcheva, T.G., 1964, Oldgaminoidnyye brakhiopody iz permi Zakavkazya. *Paleontologicheskii Zhurnal*, 1964, no. 3, 58-72. (in Russian)
- Sarytcheva, T.G. and Sokolskaya, A.N., 1952, Opredelitel paleozoyskikh brakhiopod Podmoskovnoy Kotloviny. *Tr. Paleont. Inst.*, **38**, 1-307. (in Russian)
- Sarytcheva, T.G. and Sokolskaya, A.N., 1959, O klassifikatsin lozhnoporistykh brakhiopod. *Doklady, Akad. Nauk SSSR*, **125**, 181-184. (in Russian)
- Schellwien, E., 1900, Die Fauna der Trogkofelschichten in den Karnischen Alpen und den Karawanken, 1 Teil: Die Brachiopoden, *Abh. k. k. Geol. Reichsanst.*, **16**, 1-122.
- Schréter, Z., 1963, A Bükkhegység felső-permi Brachiopodái. *Geologica Hungarica, Ser. Palaeontologica*, **28**, 79-179.
- Schuchert, C., 1929, Classification of brachiopod genera, fossil and recent. In Schuchert, C. and Le Vene, C.M., *Fossilium Catalogus, vol. 1. Animalia*. W. Junk, Berlin, 10-25.
- Shi, G.R., Archbold, N.W. and Zhan, L., 1995, Distribution and characteristics of mixed (transitional) mid-Permian (Late Artinskian-Ufimian) marine faunas in Asia and their palaeogeographical implications. *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, **114**, 241-271.
- Shimizu, D., 1961, Brachiopod fossils from the Permian Maizuru Group. *Mem. Coll. Sci., Univ. Kyoto, Ser. B*, **27**, 309-350.
- Simic, V., 1933, Gornji Perm u zapadnoj Sibiji: Das Oberperm in Westserbien. *Raspr. Geol. Inst. Kralj. Jugoslavije*, **1**, 1-130.
- Sremac, J., 1986, Middle Permian brachiopods from the Velebit Mts. (Croatia, Yugoslavia). *Palaeontologia Jugoslavica*, **35**, 1-43.
- Stehli, F.G., 1954, Lower Leonardian Brachiopoda of the Sierra Diablo. *Bull. Amer. Mus. Nat. Hist.*, **105**, 263-358.
- Stepanov, D.L., 1937, Permskie brakhiopody Spitsbergena. *Tr. Arkt. Inst.*, **76**, 105-192. (in Russian)
- Tazawa, J., 1974, *Waagenoconcha* (Brachiopoda) from the Permian of the southern Kitakami Mountains, northeast Japan. *Jour. Fac. Sci., Hokkaido Univ., Ser. 4*, **16**, 121-143.

- Tazawa, J., 1976, The Permian of Kesenuma, Kitakami Mountains: A preliminary report. *Earth Science (Chikyu Kagaku)*, **30**, 175-185.
- Tazawa, J., 1979, Middle Permian brachiopods from Matsukawa, Kesenuma region, southern Kitakami Mountains. *Saito Ho-on Kai Mus. Nat. Hist., Res. Bull.*, no. 47, 23-35.
- Tazawa, J., 1987, Permian brachiopod faunas of Japan and their palaeobiogeography. *Chikyu Monthly (Gekkan Chikyu)*, **9**, 252-255. (in Japanese)
- Tazawa, J., 1991, Middle Permian brachiopod biogeography of Japan and adjacent regions in east Asia. In Ishii, K., Liu, X., Ichikawa, K. and Huang, B., eds., *Pre-Jurassic geology of Inner Mongolia, China: Report of China-Japan Cooperative Research Group, 1987-1989*, Matsuya Insatsu, Osaka, 213-230.
- Tazawa, J., 1998, Pre-Neogene tectonic divisions and Middle Permian brachiopod faunal provinces of Japan. *Proc. Roy. Soc. Vict.*, **110**, 281-288.
- Tazawa, J., 1999a, *Leptodus* and *Spiriferella* (Permian Brachiopoda) from the Usuginu Conglomerate, southern Kitakami Mountains, northeast Japan. *Sci. Rep., Niigata Univ., Ser. E*, no. 14, 1-13.
- Tazawa, J., 1999b, Boreal-type brachiopod *Yakovlevia* from the Middle Permian of Japan. *Paleontological Research*, **3**, 88-94.
- Tazawa, J. and Gunji, Y., 1982, Middle Permian brachiopods from the Oashi Formation, Abukuma Mountains, northeast Japan. *Saito Ho-on Kai Mus. Nat. Hist., Res. Bull.*, no. 50, 67-75.
- Tazawa, J. and Matsumoto, T., 1998, Middle Permian brachiopods from the Oguradani Formation, Ise district, Hida Gaiken Belt, central Japan. *Sci. Rep., Niigata Univ., Ser. E*, no. 13, 1-19.
- Tazawa, J., Ono, T. and Hori, M., 1998, Two Permian lytoniid brachiopods from Akasaka, central Japan. *Paleontological Research*, **2**, 239-245.
- Tazawa, J., Takizawa, F. and Kamada, K., 2000, A Middle Permian Boreal-Tethyan mixed brachiopod fauna from Yakejima, southern Kitakami Mountains, NE Japan. *Sci. Rep., Niigata Univ., Ser. E*, no. 15, 1-21.
- Tazawa, J., Yoshida, K. and Machiyaama, H., 1999, Permian brachiopods from a sandstone block in the Usuginu Conglomerate, southern Kitakami Mountains, northeast Japan and their geological significance. *Earth Science (Chikyu Kagaku)*, **53**, 159-163. (in Japanese)
- Termier, H. and Termier, G., 1960, Contribution á la classification des Brachiopodes: le lophophore des Collophorides nov. ord.; Appendice. Les Oldhaminidés du Cambodge. *Soc. Géol. France, Bull., Sér. 7*, **1**, 233-243.
- Termier, H. and Termier, G., 1970, Les Productoides du Djoulfien (Permien supérieur) dans la Tethys orientale: essai sur l'agonie d'un phylum. *Ann. Soc. Géol. Nord.*, **90**, 443-461.
- Thomas, G.A., 1957, Oldhaminid brachiopods in the Permian of northern Australia. *Jour. Palaeont. Soc. India*, **2**, 174-182.
- Tong, Z., 1978, Phylum Brachiopoda; Carboniferous and Permian. In Geological Institute of southwest China, ed., *Palaeontological atlas of southwest China; Sichuan, pt. 2. Carboniferous to Mesozoic*, Geol. Pub. House, Beijing, 210-267. (in Chinese)
- Tschernyschew, F., 1902, Verkhnekamennougolnyya brachiopody Urala i Timana. *Tr. Geol. Kom.*, **16**, 1-749. (in Russian)
- Waagen, W., 1883, Salt Range fossils, 1. Productus-Limestone fossils. *Palaeontologia Indica, Ser. 13*, **1**, 391-546.
- Waagen, W., 1884, Salt Range fossils, 1. Productus-Limestone fossils. *Palaeontologia Indica, Ser. 13*, **1**, 547-728.
- Wang, C. and Yang, S., 1998, *Late Carboniferous-Early Permian brachiopods of central Xinjiang and their biostratigraphical studies*. Geol. Pub. House, Beijing, 156 p. (in Chinese)
- Wang, G., Liu, Q., Jin, Y., Hu, S., Liang, W. and Liao, Z., 1982, Phylum Brachiopoda. In

- Nanjing Institute of Geology and Mineral Resources, ed., *Palaeontological atlas of east China*, Geol. Pub. House, Beijing, 186-256. (in Chinese)
- Wang, S., 1984, Phylum Brachiopoda. In Regional Geological Surveying Team of Hubei, ed., *The palaeontological atlas of Hubei Province*, Hubei Sci. Tech. Press, Wuhan, 128-235. (in Chinese)
- Wang, Y., Jin, Y. and Fang, D., 1964, *Brachiopod fossils of China, pt. 1*. Sci. Press, Beijing, 354 p. (in Chinese)
- Wanner, J. and Sieverts, H., 1935, Zur Kenntnis der permischen Brachiopoden von Timor, 1. Lytoniidae und ihre biologische und stammesgeschichtliche Bedeutung. *Neu. Jahr. Min. Geol. Palaeont., Ser. B*, **74**, 201-281.
- Waterhouse, J.B., 1968, The classification and descriptions of Permian Spiriferida (Brachiopoda) from New Zealand. *Palaeontographica, Abt. A*, **129**, 1-94.
- Waterhouse, J.B., 1975, New Permian and Triassic brachiopod taxa. *Pap., Dep. Geol. Univ. Qd.*, **7**, 1-23.
- Waterhouse, J.B. and Waddington, J., 1982, Systematic descriptions, paleoecology and correlations of the Late Paleozoic subfamily Spiriferellinae (Brachiopoda) from the Yukon Territory and the Canadian Arctic Archipelago. *Geol. Soc. Canada, Bull.*, **289**, 1-73.
- Waterhouse, J.B., Waddington, J.W. and Archbold, N., 1978, The evolution of the Middle Carboniferous to the Late Permian brachiopod subfamily Spiriferellinae Waterhouse. In Stelck, C.R. and Chatterton, B.D.E., eds., *Western and Arctic Canadian biostratigraphy. Geol. Assoc. Canad., Spec. Pap.*, **18**, 415-443.
- Williams, A., Harper, D.A.T. and Grant, R.E., 2000, Lytoniida. In Williams, A., Brunton, C.H.C. and Carlson, S.J., eds., *Treatise on Invertebrate Paleontology, pt. H. Brachiopoda Revised, vol. 3. Linguliformea, Craniiformea, and Rhynchonelliformea (part)*, Geol. Soc. Amer., Boulder and Univ. Kansas, Lawrence, 619-642.
- Wiman, C., 1914, Über die Karbonbrachiopoden Spitzbergens und Beeren Eilands. *Nova Acta Regiae Soc. Sci. Upsaliensis, Ser. 4*, **3**, 1-91.
- Yabe, H., 1900, The brachiopod *Lyttonia* from Rikuzen Province. *Jour. Geol. Soc. Tokyo*, **7**, 1-4.
- Yanagida, J., 1963, Brachiopods from the Upper Permian Mizukoshi Formation, central Kyushu. *Mem. Fac. Sci., Kyushu Univ., Ser. D*, **14**, 69-78.
- Yanagida, J., 1996, Permian brachiopods from the Tsunemori Formation, SW Japan, and their paleobiogeographic implication. In Copper, P. and Jin, J., eds., *Brachiopods*, A.A. Balkema, Rotterdam, 313-315.
- Yang, D., 1984, Systematic description of palaeontology: Brachiopoda. In Yichang Institute of Geology and Mineral resources, ed., *Biostratigraphy of the Yangtze Gorge area, (3) Late Palaeozoic Era*, Geol. Pub. House, Beijing, 203-239, 330-333, 387-396. (in Chinese)
- Yang, D., Ni, S., Chang, M. and Zhao, R., 1977, Phylum Brachiopoda. In Hubei Institute of Geological Science et al., eds., *Palaeontological atlas of mid-south China, pt. 2. Late Palaeozoic volume*, Geol. Pub. House, Beijing, 303-470. (in Chinese)
- Zavodovsky, V.M., Stepanov, D.L., Balashova, E.A., Eltyshova, R.S., Lobanova, O.V., Lyutkevich, E.M., Miklukho-Maklay, A.D., Nekhoroshev, V.P., Popov, Yu.N., Radchenko, G.P. and Sokolov, B.S., 1970, *Polevoy atlas permskoy fauny i flory Severo-Vostoka SSSR*. Magadanskoe Knizhnoe Izdatelstvo, Magadan, 407 p. (in Russian)
- Zhan, L., 1979, Descriptions of fossils: Brachiopoda. In Hou, H., Zhan, L., Chen, B. et al., *The coal-bearing strata and fossils of Late Permian from Guangtung*, Geol. Pub. House, Beijing, 61-100. (in Chinese)
- Zhan, L. and Wu, R., 1982, Early Permian brachiopods from Xainza district, Xizang (Tibet). In CGQXP Editorial Committee, Ministry of Geology and Mineral Resources, P. R. China, ed., *Contribution to the geology of the Qinghai-Xizang (Tibet) Plateau, vol. 7*, Geol. Pub. House, Beijing, 86-109. (in Chinese)

- Zhang, Y. and Ching, Y., 1961, An Upper Permian brachiopod fauna from Jingxian, Anhui Province. *Acta Palaeont. Sinica*, **9**, 401-417. (in Chinese).
- Zhang, Y., Fu, L., Ding, P. and Qi, W., 1983, Phylum Brachiopoda. In Xian Institute of Geology and Mineral Resources, ed., *Palaeontological atlas of northwest China, Shaanxi, Gansu and Ningxia volume, pt. 2. Upper Palaeozoic*, Geol. Pub. House, Beijing, 244-425. (in Chinese)
- Zhu, T., 1990, *The Permian coal-bearing strata and palaeobiocoenosis of Fusian*. Geol. Pub. House, Beijing, 127 p. (in Chinese)

Explanation of Plates

(All figures are in natural size)

Plate 1

Figs. 1-3. *Transennatia gratiosa* (Waagen)

1. Internal mould of a ventral valve, NU-B305; 2. External latex cast of a ventral valve, NU-B302; 3. Anterior view of an internal mould of a ventral valve, NU-B304.

Fig. 4. *Waagenoconcha* sp.

External mould of a dorsal valve, NU-B336.

Fig. 5. *Tyloplecta* cf. *yangzeensis* (Chao)

External mould of a dorsal valve, NU-B335.

Figs. 6a, b. *Cancrinella* sp.

External mould and external latex cast of a ventral valve, NU-B307.

Figs. 7-10. *Leptodus nobilis* (Waagen)

Internal moulds of four ventral valves, 7. NU-B308; 8. NU-B309; 9. NU-B313; 10. NU-B314.

Figs. 11-13. *Linoproductus cora* (d'Orbigny)

11a, b. External latex cast and internal mould of a ventral valve, NU-B337; 12, 13. Internal moulds of two ventral valves, 12. NU-B357; 13. NU-B341.

Plate 1




Plate 2

Figs. 1-8. *Linoproductus cora* (d'Orbigny)

1a, b, c, d. Ventral, posterior, anterior and lateral views of an internal mould of a ventral valve, NU-B344; 2, 3. External latex casts of two dorsal valves, 2. NU-B361; 3. NU-B362; 4-6. Internal moulds of three ventral valves, 4. NU-B351; 5. NU-B352; 6. NU-B350; 7. External latex cast of a ventral valve, NU-B340; 8. External mould of a dorsal valve, NU-B360.

Figs. 9a, b. *Derbyia grandis* Waagen

External latex cast and internal mould of a ventral valve, NU-B364.

Plate 2




Plate 3

Figs. 1, 2, 4. *Derbyia nipponica* Nakamura

1a, b. External latex cast and internal mould of a ventral valve, NU-B365; 2. Internal mould of a tral valve, NU-B367.4. Internal mould of a ventral valve, NU-B367.

Figs. 3a-c. *Derbyia grandis* Waagen

External latex cast, internal mould and internal latex cast of a ventral valve, NU-B363.


Plate 3




1a



2



3a




3b



1b



4



3c

Plate 4

Figs. 1-10. *Spiriferella keilhavii* (von Buch)

1a, b. Ventral and dorsal views of internal mould of a conjoined valve, NU-B325; 2. External latex cast of a ventral valve, NU-B317; 3. External latex cast of a dorsal valve, NU-B321; 4. External latex cast of a ventral valve, NU-B318; 5a, b. Internal mould and internal latex cast of a dorsal valve, NU-B332; 6. Internal mould of a ventral valve, NU-B326; 7a, b. External latex cast and internal mould of a ventral valve, NU-B315; 8a, b. Internal mould and internal latex cast of a ventral valve, NU-B329; 9a, b. Internal mould and internal latex cast of a dorsal valve, NU-B334; 10a, b. Internal mould and internal latex cast of a ventral valve, NU-B327.


Plate 4



1a




1b



2



3




4




5a



6




7a



5b




7b




8a




8b



9a



9b



10b



10a