

***Isogramma* (Brachiopoda) from the lower Copacabana Group of the Lake Titicaca region, Bolivia**

Masayuki FUJIKAWA*, Takeshi ISHIBASHI** and Jun-ichi TAZAWA***

Abstract

A Late Carboniferous brachiopod *Isogramma paotchowensis* (Grabau and Chao, 1928) is described for the first time from the lower Copacabana Group of Yampupata in the Lake Titicaca region, Bolivia. The occurrence of *I. paotchowensis* suggests that the fossil horizon Yp-26 of the lower Copacabana Group is assigned to Late Carboniferous in age.

Key words: Bolivia, brachiopod, lower Copacabana Group, *Isogramma paotchowensis*, Lake Titicaca region.

Introduction

The brachiopod specimens described in this paper were collected by two of the authors (MF and TI) in September 1998, from tuffaceous sandy shale of the lower Copacabana Group at Yampupata in the Lake Titicaca region, Bolivia, nearby the border with Peru (Fig. 1). The Devonian to Permian formations including the Copacabana Group are exposed in this area. Late Carboniferous and Lower Permian rich faunas are known from the Copacabana Group since d'Orbigny's (1842) work in Bolivia and Central Andes. Many paleontological papers have been published by Newell (1949), Urdininea and Yamagiwa (1980), Arelleno (1983), Maeda and Sakagami (1983), Nagai (1983), Sakagami (1984), Kobayashi and Hamada (1986), Suarez-Riglos et al. (1987), Wilson (1990), Sakagami and Mizuno (1994) and Sakagami (1995a, 1995b). The brachiopod fauna has also been reported by Kozłowski (1914), Chronic (in Newell, 1949), Newell et al. (1953), Branisa (1965), Samtleben (1971), Yanagida (1983), and Sakagami (1984). Recently, Ishibashi and Fujikawa (1999) reported the following

* Graduate School of Science and Technology, Niigata University, Niigata 950-2181, Japan

** Kashii-dai 2-14-10, Fukuoka 813-0014, Japan

*** Department of Geology, Faculty of Science, Niigata University, Niigata 950-2181, Japan

(Manuscript received 20 January, 2003; accepted 17 February, 2003)




Fig. 1. Index map showing the fossil locality of the Yampupata area in the Lake Titicaca region, Bolivia.

four brachiopod species from the Yampupata area: *Linoproductus* sp., *Isogramma* sp., *Stereochia* sp., and *Neospirifer* sp.

The purpose of this paper is to describe a brachiopod species *Isogramma paotchowensis* (Grabau and Chao, 1928) from the Yampupata area, and to suppose the age of the lower Copacabana Group in this area. Among the authors, MF and JT are responsible for the systematic descriptions, and MF and TI are responsible for the field geology. All the specimens of *I. paotchowensis* are housed in the Department of Geology, Faculty of Science, Niigata University.

Stratigraphy

The Copacabana Group is narrowly distributed in Peru and Bolivia through the Andes, with the strike of NW-SE. This group was recognized by Newell et al. (1953) as the Upper Carboniferous and Lower Permian, consisting of massive limestone and black shale, about 1900 m in the maximum thickness.

The geological map of the Yampupata area has been made by the Geological Survey of Bolivia (GEOBOL) (1978). In the Yampupata area, the Copacabana Group overlies the Devonian Aigachi Formation and the Carboniferous Khasa and Cumana formations, and is covered with unconformity by the Tertiary rocks. The lower part of the Copacabana Group, having a general trend NW-SE, and dipping 40° NE, is estimated more than 340 m in thickness




Fig. 2. Columnar section of the lower Copacabana Group at Yampupata, showing the fossil horizon Yp-26, revised from Sakagami (1986).



Fig. 3. Outcrop of the lower Copacabana Group (Yp-26) at Yampupata.

(Sakagami, 1986). Fig. 2 shows the columnar section of the lower Copacabana Group at Yampupata. We have collected the brachiopod specimens from pale green tuffaceous sandy shale at the horizon Yp-26 of Sakagami et al. (1985), 130 m above from the base of the Copacabana Group (Fig. 3). Bryozoans, brachiopods, cephalopods, bivalves, and gastropods have been reported from this bed (Sakagami, 1986).

In this area, the Carboniferous–Permian boundary has not settled the precise horizon yet. Sakagami (1986) mentioned that the horizon Yp-26 lies on the 80m above the Carboniferous–Permian boundary. Suarez-Riglos et al. (1987) studied the conodont biostratigraphy of the Copacabana Group in Bolivia, and they proposed that the lowermost Permian *Triticites* Zone and *Triticites nitens* Subzone are really equivalent to the uppermost Carboniferous. This proposal was followed by Sakagami et al. (1991) who examined along some routes, Cuyavi, Yampupata, Ancoraimes, and Matilde, in the Lake Titicaca region. Sakagami and Mizuno (1994) discovered the Middle Pennsylvanian fusulinaceans and conodonts from the fossil horizons Cu-04, 11, 29, 32, 39, 43, 46a, and 46b of the Cuyabi route (Sakagami, 1986) and Yp-10 of the Yampupata route in the lower part of the Copacabana Group. The horizon Cu-46 in the Cuyabi route were referable to Yp-10 (about 45 m below Yp-26) in the Yampupata route from the fusulinacean evidence (Sakagami and Mizuno, 1994).

Consequently, the horizon Yp-26 is referable to Middle Pennsylvanian or later. Meanwhile, *Isogramma paotchowensis*, which was collected at the horizon Yp-26, has only occurred from the Upper Carboniferous of north China and the Carnic Alps. Therefore, we correlate the sampling horizon Yp-26 of the lower Copacabana Group with Late Carboniferous in age. This view does not conflict with the previous study of Sakagami and Mizuno (1994), and is valuable for the argument of the Carboniferous–Permian boundary in the lower Copacabana Group of the Yampupata area.

Systematic paleontology

Order Dictyonellida Cooper, 1956
 Superfamily Eichwaldioidea Schuchert, 1893
 Family Isogrammidae Schuchert, 1929
 Genus *Isogramma* Meek and Worthen, 1870

Isogramma paotchowensis (Grabau and Chao, 1928)

Fig. 4.

Aulacorhynchus paotchowensis Grabau and Chao, in Chao, 1928, p. 33, pl. 1, fig. 27; pl. 4, figs. 1-5.

Isogramma paotchowensis (Grabau and Chao). Aigner and Heritsch, 1931, p. 307, pl. 2, figs. 29-36; pl. 3, figs. 37-44; pl. 4, figs. 45-51; pl. 5, figs. 52, 53, 67-71; Hatai and Omura,




Fig. 4. *Isogramma paotechowensis* (Grabau and Chao, 1928), from the lower Copacabana Group of Yampupata in the Lake Titicaca region, Bolivia. 1: ventral valve, NU-B591. 2: dorsal valve, NU-B593. 3: ventral valve, NU-B592. 4: dorsal valve, NU-B594.

1941, pl. 2, fig. 4; Wang, 1957, p. 160, pl. 92, fig. 14; Wang et al., 1964, p. 354, pl. 37, fig. 38; Lee and Duan, 1985, p. 239, pl. 66, figs. 14, 15, 23.

Isogramma paotechowense (Grabau and Chao). Licharev, in Gorsky, 1939, p. 83, pl. 17, figs. 1, 2.

Isogramma sp. Ishibashi and Fujikawa, 1999, pl., figs. 4-8.

Material.—Six specimens: two ventral valves (NU-B591, NU-B592), a single dorsal valve (NU-B593), and three valve fragments (NU-B594-NU-B596).

Description.—Shell medium size for genus, transversely suboval in outline, with greatest width slightly anterior to hingeline; length 44 mm, width 72 mm in the largest specimen (NU-B591). Ventral valve slightly convex. Dorsal valve slightly concave to almost flattened. External surface of both valves ornamented by fine numerous and regular concentric fila. Fila sharply separated by shallow grooves; numbering 9-12 in 5 mm at mid of ventral valve.

Remarks.—The specimens from Yampupata, Lake Titicaca region have been figured and reported as *Isogramma* sp. by Ishibashi and Fujikawa (1999) without any systematic description. These specimens are identical with *Isogramma paotechowensis* (Grabau and Chao, 1928), originally described by Grabau and Chao (in Chao, 1928) as *Aulacorhynchus paotechowensis*

from the Taiyuan Series of Shanxi, north China in size, shape and external ornament of shell. The number of fila is 10 per 5 mm in the Chinese type specimen (Chao, 1928, pl. 1, fig. 27), and 9-12 per 5 mm in the Bolivian specimens.

The shells, described and figured as *Isogramma paotechowensis*, from the Upper Carboniferous and Lower Permian of Fergana (Volgin, 1957, p. 39, pl. 1, figs. 8-9; Volgin, 1960, p. 41, pl. 2, fig. 6; Sergunkova and Zhizhilo, 1974, p. 68, pl. 12, figs. 1-3), and from the Middle Permian of the southern Kitakami Mountains, northeast Japan (Minato, 1955, p. 29, text-fig. 1; Nakamura, 1970, p. 306, pl. 3, figs. 1, 2; pl. 4, figs. 1, 2), may be not assigned to *I. paotechowensis* in their much larger size and more transverse outline.

Distribution. — Upper Carboniferous of the Carnic Alps; Shaanxi, Shanxi and Hebei, north China; Lake Titicaca region, Bolivia.

Acknowledgments

We would like to express our gratitude to Dr. O. Sanjines of Servicio Geologico de Bolivia (GEOBOL) for his kind offer of the local information. Thanks are also expressed to Prof. R. Santivanez of Universidad Mayor de San Andres for his sincere cooperation in the field work. Our sincere thanks are due to Dr. I. Niikawa of Niigata University for critical reading the manuscript. This study was financially supported in part by a grant from the Association of Geology in Japan in 1998.

References

- Aigner, G. and Heritsch, F., 1931, Das Genus *Isogramma* im Carbon der Südalpen. *Denkschr. Akad. Wiss. Wien, Math. Naturwiss. Kl.*, **102**, 303-316.
- Arellano, J., 1983, Trilobites del Permico Inferior de Bolivia. *Bull. Inst. Frances de Estudios Andinos, Lima, Peru*, **12**, 91-102.
- Branisa, L., 1965, Los Fosiles guias de Bolivia, 1. Paleozoico. *Ser. Geol. Bolivia*, **6**, 1-282.
- Chao, Y.T., 1928, Productidae of China, Part 2, Chonetinae, Productinae and Richthofeninae. *Palaeont. Sin., Ser. B.*, **5**, fasc. 3, 1-103.
- Cooper, G.A., 1956, Chazy and related brachiopods. *Smithson. Misc. Coll.*, **127**, 1-1245.
- GEOBOL (Servicio Geologico de Bolivia), 1978, *Mapa geologico de Bolivia, Memoria explicativa. Escala 1:1,000,000*. La Paz, Bolivia.
- Gorsky, I.I., 1939, *Atras rukovodyaschikh form iskopaemykh faun SSSR*, **5**, *Spedny i verkhniy otdely kamennougolnoy sistemy*. GONTI, Leningrad, 179 p. (in Russian)
- Hatai, K. and Omura, T., 1941, On a species of *Isogramma* (Brachiopoda) from Manchoukuo. *Jour. Geol. Soc. Japan*, **48**, 43-46.
- Ishibashi, T. and Fujikawa, M., 1999, Permian fauna of the Copacabana Group distributed around the Lake Titicaca in the Central Andes. *Jour. Geogr.*, **108**, no. 3, 321-327.
- Kobayashi, T. and Hamada, T., 1986, A new Permian genus of Trilobita from Bolivia. *Proc. Japan Acad., Ser. B*, **62**, 181-183.
- Kozłowski, R., 1914, Les brachiopodes du Carbonifère supérieur de Bolivie. *Ann. Paléont.*, **9**, 1-100.

- Lee, L. and Duan, C., 1985, Brachiopoda, *In* Paleontological Atlas of North China, (1) Paleozoic Volume, *Bull. Tianjin Inst. Geol. Min. Res.*, no. 10, 209-260. (in Chinese)
- Maeda, S. and Sakagami, S. (eds.), 1983, On the Geology and Paleontology in the Andes. *Jour. Geogr.*, **92**, 1-54. (in Japanese)
- Meek, F.B. and Worthen, A.H., 1870, Descriptions of new species and genera of fossils from the Paleozoic rocks of the western states. *Acad. Nat. Sci. Phila., Proceed., Ser. 2*, **22**, 22-56.
- Minato, M., 1955, *Isogramma paotechowensis* (Grabau and Chao) from the Permian of Japan. *Trans. Proc. Palaeont. Soc. Japan, N. S.*, no. 18, 29-30.
- Nagai, K., 1983, On the sedimentary facies of the Copacabana Group (Permian). *Jour. Geogr.*, **92**, 45-48. (in Japanese)
- Nakamura, K., 1970, *Isogramma* from the Permian Kanokura Series of the Kitakami Mountains, Japan. *Jour. Fac. Sci., Hokkaido Univ., Ser. 4*, **14**, 301-311.
- Newell, N.D., 1949, Geology of the Lake Titicaca region, Peru and Bolivia. *Geol. Soc. Amer., Mem.* **36**, 111 p.
- Newell, N.D., Chronic, J. and Roberts, T.G., 1953, Upper Paleozoic of Peru. *Geol. Soc. Amer., Mem.* **58**, 276 p.
- Orbigny, A. d', 1842, *Voyages dans l'Amerique Meridionale de 1826-1833, Paléontologie*. Pitois-Levrault et Cie, Paris, 188 p.
- Sakagami, S., ed., 1984, *Biostratigraphic study of Paleozoic and Mesozoic Groups in Central Andes - An interim report -*. Dept. Earth Sci., Fac. Sci., Chiba Univ., 82 p.
- Sakagami, S., ed., 1986, *Biostratigraphic study of Paleozoic and Mesozoic Groups in Central Andes. - An interim report (2) -*. Dept. Earth Sci., Fac. Sci., Chiba Univ., 83 p.
- Sakagami, S., 1995a, Upper Paleozoic bryozoans from the Lake Titicaca region, Bolivia, Pt. 1, Introductory remarks, stratigraphy and systematic paleontology. *Trans. Proc. Palaeont. Soc. Japan, N. S.*, no. 180, 226-260.
- Sakagami, S., 1995b, Upper Paleozoic bryozoans from the Lake Titicaca region, Bolivia, Pt. 2, Systematic paleontology. *Trans. Proc. Palaeont. Soc. Japan, N. S.*, no. 180, 261-281.
- Sakagami, S., Kawabe, T., Ishibashi, T., Nagai, K., Carrasco, R., Escobar, A., and Rangel, C., 1985, Biostratigraphic study of Paleozoic and Mesozoic Groups in Central Andes (3) – Preliminary Report of the Field Work in 1984-. *Jour. Geogr.*, **94**, 34-44. (in Japanese)
- Sakagami, S. and Mizuno, Y., 1994, Discovery of Middle Pennsylvanian fusulinaceans and conodonts from the Copacabana Group in the Lake Titicaca region, Bolivia. *Trans. Proc. Palaeont. Soc. Japan, N. S.*, no. 174, 484-494.
- Sakagami, S., Yanagida, J., Kawabe, T., Mizuno, Y., Okamoto, K., Garcia, R., Rangel, C., and Aldana, M., 1991, Paleontological analysis on Late Paleozoic Era in Central Andes –Report on Field Work in 1990-. *Jour. Geogr.*, **100**, 433-443. (in Japanese)
- Samtleben, C., 1971, Zur Kenntnis der Productiden und Spiriferiden des bolivianischen Unterperms. *Reih. Geol. Jahrb.*, **3**, 3-138.
- Schuchert, C., 1893, Classification of the Brachiopoda. *American Geologist*, **11**, 141-167.
- Schuchert, C., 1929, Classification of brachiopod genera, fossil and recent. *In* Schuchert, C. and LeVene, C.M., *Animalia pars 42. In Pompeckj, J.F., ed., Fossilium Catalogus*, **1**, Junk. Berlin, 10-25.
- Sergunkova, O.I. and Zhizhilo, O.R., 1974, Brachiopody srednego karbona, verkhnego karbona i nizhney permi Fergany. *In* Khodanovich, R.L., ed., *Biostratigrafiya verkhnego paleozoya gornogo obramleniya Yuzhnoi Fergany*. Izd-vo "Fan" uz SSR, Tashkent, 54-77. (in Russian)
- Suarez-Riglos, M., Hunicken, M.A., and Merino, D., 1987, Conodont biostratigraphy of the Upper Carboniferous-Lower Permian rocks of Bolivia. *In* Austin, R., ed., *Conodont -Investigative techniques and applications-*. Ellis Horwood, 422 p.
- Urdininea, M. and Yamagiwa, N., 1980, Paleontological study on the Copacabana Group at

- the hill of Jacha Khatawi in the Yaurichambi area, Bolivia, South America, Part 1, Fusulinids. *Prof. Saburo Kanno Mem. Vol.*, 277-289.
- Volgin, V. I., 1957, Izogrammidy Yuzhnoy Fergany. *Vestnik Leningradskogo Univ.*, no. 18, 34-42. (in Russian)
- Volgin, V. I., 1960, Brakhiopody verkhnekamennougolnykh i nizhnepersmskikh otlozheniy Yuzhnoy Fergany. *Izdatelstvo Leningradskogo Univ.*, 203 p. (in Russian)
- Wang, Y., 1957, Phylum Brachiopoda. In Yang, K. and Wang, Y., *Index fossils of China, Invertebrate, Part 2*, Geological Publishing House, Beijing, 109-171. (in Chinese)
- Wang, Y., Jin, Y. and Fang, D., 1964, *Brachiopod fossils of China, Part 1*. Science Press, Beijing, 354 p. (in Chinese)
- Willson, E. C., 1990, Permian corals of Bolivia. *Jour. Paleont.*, **64**, 60-78.
- Yanagida, J., 1983, Lower Permian brachiopods from Peru and Bolivia. *Jour. Geogr.*, **92**, 51-54. (in Japanese)