

## ***Monodiexodina* from the Permian Oguradani Formation, Hida Gaaien Belt, central Japan**

Katsumi UENO\* and Jun-ichi TAZAWA\*\*

### **Abstract**

A well-known antitropical fusulinoidean genus *Monodiexodina* is reported with illustration of specimen for the first time from calcareous sandstone of the Oguradani Formation in the Ise area, Hida Gaaien Belt, central Japan. The present Oguradani specimens are referable to *Monodiexodina* sp., and indicate a middle Middle Permian (Murgabian/Wordian) age. The occurrence of *Monodiexodina* from the Permian Oguradani Formation, together with the overall similarity of Paleozoic stratigraphy in the two belts, further reinforces the current opinion on the original geologic connection of the Hida Gaaien Belt with the South Kitakami Belt of Northeast Japan.

*Key words:* central Japan, Fusulinoidea, Hida Gaaien Belt, Middle Permian, *Monodiexodina*, Oguradani Formation.

### **Introduction**

The Hida Gaaien Belt in central Japan is a unique geotectonic domain composed mainly of Ordovician mafic- and ultramafic rocks, about 300-400Ma high-P/T metamorphic rocks, and Paleozoic and Mesozoic non-metamorphosed rocks including Permo-Carboniferous shallow-marine strata, and is narrowly distributed along the margin of the Hida Belt (Tazawa, 1989). The Permian in the Hida Gaaien Belt is distributed, from east to west, in the Fukuji, Moribu, and Ise areas (Fujimoto et al., 1962; Igo, 1990). Among them, the Mizuyagadani Formation in the Fukuji area and the Moribu Formation in the Moribu area have been studied stratigraphically and paleontologically in detail, and correlated to the Early Permian (Igo, 1957; Niikawa, 1980; Niko et al., 1987) and the late Early to Middle (and partly Late?) Permian

---

\* Department of Earth System Science, Faculty of Science, Fukuoka University, Fukuoka 814-0180, Japan

\*\* Department of Geology, Faculty of Science, Niigata University, Niigata 950-2181, Japan  
(Manuscript received 5 February, 2004 accepted 25 February, 2004)

(Yamada and Yamano, 1980; Horikoshi et al., 1987; Tazawa et al., 1993; Tazawa, 1996a, 2001), respectively. In contrast with these formations, the Permian of the Ise area in the western part of the Hida Gaien Belt still remains being poor in information on its stratigraphy and fossil contents.


In the Ise area, several stratigraphic units such as the Oguradani, Konogidani, Magatoji, Oboradani, and Otani formations have been proposed for the non-metamorphosed Permian (Ozaki et al., 1954; Yamada et al., 1958; Yamada, 1967). Yamada (1967) once mentioned the occurrence of *Monodiexodina* sp. from calcareous shale of the Oguradani Formation but without illustration of specimen. With respect to this Oguradani *Monodiexodina*, Tazawa et al. (1993) noted, based on personal communication with K. Yamada, that the fusulinoidean specimens of Yamada (1967) have been lost already and, moreover, additional collection of specimens from the same locality would be very difficult due to submergence by the subsequent construction of the Kuzuryu Dam (Kuzuryu Lake). *Monodiexodina* is an important fusulinoidean genus for the discussion on the origin and paleobiogeographic affinity of the Hida Gaien Belt, and thus additional specimens from the Oguradani Formation is eager to be reported and illustrated.

In this paper, we report the mode of occurrence of *Monodiexodina* from the Oguradani Formation and illustrate the specimen, although they are by no means well preserved. We further refer to some geologic implications of the occurrence of *Monodiexodina* from the formation. In the Hida Gaien Belt, this genus has only hitherto been reported by Tazawa et al. (1993) with illustration of specimen from sandstone of the middle part of the Moribu Formation. Therefore, this is the second report of occurrence of *Monodiexodina* with illustration of specimen from the Hida Gaien Belt. Specimens examined in this study are all registered and housed in the paleontological collections of the Department of Geology, Faculty of Science, Niigata University, Niigata, Japan, with prefix NU-F.

### Stratigraphy

The Oguradani Formation is mainly distributed as an E-W trending narrow belt from Kuzuryu Lake to Motodo through Shiroki-yama, and is fine clastic-dominated strata with minor intercalations of limestone. It was first introduced by Ozaki et al. (1954) as one of Permian stratigraphic units in the Ise area. Later, Kawai et al. (1957), in the explanatory text of 1:50,000 geologic map "Arashimadake" sheet, regarded it as a member in the Nojiri Formation.

Yamada (1967) summarized the Permian of the Ise area, and recognized two formations in the Nojiri Group, the Oguradani Formation in apparently lower stratigraphic position and the Konogidani Formation in upper stratigraphic position. He noted that the Oguradani Formation in the type section at Oguradani is of sandstone-dominated in the lower one third and of shale-dominated in the upper two-thirds. Recently, Tazawa and Matsumoto (1998) redefined the Oguradani Formation as being composed of the Oguradani and Ashidani formations of Yamada (1967), the latter of which had been treated in Yamada (1967) as a formation of



**Fig. 1.** Map showing fossil locality, HIF1 (asterisk), using topographic map of “Echizen-asahi” scale 1:25,000 published by Geographical Survey Institute of Japan.

unknown age. They noted that the formation, attaining over 750 m in total thickness, is divided lithologically into three members: the lower shale-limestone member of 470 m thick, the middle sandstone member of 110 m thick, and the upper shale member of more than 170 m thick.


Fossils occur rather scarcely in the Oguradani Formation, but such brachiopods and cephalopods as *Lyttonia* (= *Leptodus*), *Camarophoria*, *Productus*, *Paraceltites*, and *Foordiceras* have been reported by Hayasaka and Matsuo (1951) and Hayasaka and Ozaki (1955) from the formation. Recently, Tazawa and Matsumoto (1998) described a Middle Permian brachiopod fauna composed of *Derbyia* cf. *buchi* (d'Orbigny), *Meekella* sp., *Transennatia gratiosa* (Waagen), *Leptodus nobilis* (Waagen), *Stenosisma margaritovi* (Tschernyschew), and *Hustedia* cf. *grandicosta* (Davidson) from argillaceous impure limestone in the lower part of the Oguradani Formation. They concluded that the Oguradani brachiopod fauna consists of mixed Boreal-, Tethyan-, and bipolar-type genera, and can be correlated with those from the Southern Subzone of the Inner Mongolian-Japanese Transition Zone of Tazawa (1991) in Middle Permian paleobiogeography.

The present fusulinoidea-bearing samples studied herein were collected by the junior author (JT) from the locality HIF1, an isolated outcrop on an E-W trending ridge about 500 m north of the mouth of Tsuratani, north of Tomedoro (Tometoro), Izumi-mura, Ono-gun, Fukui Prefecture, i.e., the Ise area, western part of the Hida Gaien Belt, central Japan (Fig. 1). At the fossil locality, gray to greenish gray tuffaceous and calcareous sandstones are exposed on a small road cutting. The fossil-bearing sandstone probably belongs to the lower part of the Oguradani Formation (see Tazawa and Matsumoto, 1998, fig. 2), although the exact stratigraphic position has not been settled yet.

### Mode of occurrence

We have examined several rock samples from the fossil locality described in the precedent chapter. Fusulinoideans are generally contained in gray to greenish gray, medium-grained, calcareous sandstone. In most rock samples collected, however, calcareous materials including fusulinoidean shells have been completely dissolved by weathering. Consequently, these fusulinoideans exhibit typical “Matsubaishi-type” mode of occurrence (Figs. 2A-D), which has been reported as a typical mode of occurrence of *Monodioxodina matsubaishi* (Fujimoto) and its allied form in the South Kitakami Belt and the Hida Gaien Belt (Fujimoto, 1956; Tazawa et al., 1993). Fusulinoidean shells are swarmed in several particular layers (Fig. 2A), arranged almost parallel to the bedding plane, and tend to be roughly oriented (Figs. 2A, 2B, 2D). Their original external shell morphology including shell length and shell width seems to be relatively well preserved as external molds in most specimens on the surfaces almost parallel to the bedding plane even though they may have slight deformation. In contrast, fusulinoidean shells are very much squashed by compaction in sections vertical to the bedding plane (Fig. 2C).


We also collected some rock samples in which calcareous materials including fusulinoidean shell are still preserved, and processed them for thin sections (Figs. 3A-H). In this type of samples, however, fusulinoidean shells are more poorly preserved than those found in the rock samples showing the “Matsubaishi-type” mode of occurrence due to strong abrasion and subsequent diagenetic dissolution of the outer part of shells. Many fusulinoidean shells in this type of samples have irregular shell outlines surrounded by thin, dark residual materials (e.g., Fig. 3H). They always have stylolitic contact with the surrounding sediments, which consist of medium-grained calcareous sandstone (e.g., Fig. 3G). Under the microscope, a large number of angular to sub-angular detrital quartz grains cemented by calcareous matrix are observed among fusulinoidean shells. Besides fusulinoideans, other calcareous particles such as crinoid stems and fragments of bryozoans are found in thin sections. Similarly to the first type of samples, fusulinoidean shells are also very much squashed by compaction in sections perpendicular to the bedding plane.



**Fig. 2.** Photographs of *Monodiexodina*-bearing sandstone. A: *Monodiexodina*-swarmed surface almost parallel to bedding plane. Fusulinoideans (*Monodiexodina* sp.) are completely dissolved and exhibit typical “Matsubaishi-type” mode of occurrence. NU-F41,  $\times 1$ . B: Surface almost parallel to bedding plane. Molds of shells of *Monodiexodina* sp. are generally preserved in medium-grained sandstone and sometimes filled partly with brown-colored residue. NU-F42,  $\times 1$ . C: Surface almost perpendicular to bedding plane, showing linear arrangement of highly squashed individuals of *Monodiexodina* sp. Almost vertical to surface of B,  $\times 1$ . D: Several complete external molds of shells of *Monodiexodina* sp. on surface almost parallel to bedding plane. They have very elongated subcylindrical shells. NU-F43,  $\times 2$ .

### Shell morphology

In this study, we unfortunately could not get any completely oriented (axial and/or sagittal) section, which is generally indispensable for the taxonomic study of fusulinoidean foraminifers. Specimens available for this study are external molds of whole fusulinoidean shells preserved in weathered sandstone (Figs. 2A-H) and a number of randomly oriented fragments in thin



**Fig. 3.** Photomicrographs of *Monodiexodina*-bearing calcareous sandstone. A: Microfacies of calcareous sandstone (partly sandy limestone) with many, poorly preserved fusulinoidean (*Monodiexodina* sp.) shells. NU-F71,  $\times 5$ . B-H, Various fragmented and oriented specimens of *Monodiexodina* sp. B,C: Parallel sections with several volutions, in which a number of septa can be seen in one volution. NU-F60 and NU-F65, both  $\times 10$ . D-G: Various fragments of tangential and nearly tangential sections, showing an elongated subcylindrical shell, regularly fluted septa, and well-developed axial fillings. NU-F68a, NU-F68b, NU-F51a and NU-F52, all  $\times 10$ . H: Fragment of tangential section corresponding almost to a half of whole shell. Almost plane septa in central part of shell (right-hand side), regularly arranged septal loops, and heavy axial fillings in axial part are observed. This specimen should have a very elongated subcylindrical shell, attaining more than 25 mm in total shell length. NU-F51b,  $\times 10$ .

sections of calcareous sandstone (Figs. 3A-H). By joining together the lines of fragmentary information from them, however, we could reconstruct basic morphology of the Oguradani specimens, which enabled us some taxonomic discussion.

The present Oguradani fusulinoideans are about 16 to 21 mm in length and about 2 mm in width in large, molded specimens. Although it is difficult in most cases to measure the exact length and width of shells, their form ratios would attain to around 8.0 to 10.0. They have very elongated subcylindrical shells (Figs. 2D, 3H) with probably 5 to 6 volutions (Figs. 3B, 3C, 3E). Septa are almost plane in the central part of shell but tend to be fluted regularly towards polar regions, forming semicircular septal loops (Figs. 3D-H). Spirotheca is completely altered by diagenesis and looks structureless in most specimens, but obscure vestige of keriothecal structure is partly discernible. Axial fillings are well developed in axial and polar regions as secondary deposits (Figs. 3E, 3H).

All these morphological features observed in the Oguradani specimens, especially the highly elongated shells, together with their typical “Matsubaishi-type” mode of occurrence, strongly suggest that they can be subsumed in the genus *Monodiexodina*. As we have got no well-oriented section available in this study, however, the Oguradani fusulinoideans should be best assigned to *Monodiexodina* sp. in the existing state of affairs. Poor preservation of our materials notwithstanding, their “Matsubaishi-type” mode of occurrence somewhat reminds us of possibility that the present specimens may be referable to *Monodiexodina matsubaishi* (Fujimoto).

### Geologic implications

*Monodiexodina* is well-known as an antitropical fusulinoidean genus, which is mainly distributed in the northern and southern transitional zones in the context of Permian paleobiogeography (e.g., Shi et al., 1995; Ueno, 2003). Recently, Ueno and Tazawa (2003) summarized the geographic and stratigraphic distributions of *Monodiexodina*, and concluded that the genus has a range from the Kubergandian to Midian (namely the entire Middle Permian) in the northern transitional zone, i.e., the Inner Mongolian-Japanese Transition Zone of Tazawa (1991) which corresponds to present-day Northeast China, South Primorye, and several pre-Jurassic terranes in Japan including the Hida Gaien Belt. Furthermore, they noted that it is especially dominated in the Murgabian (middle Middle Permian) in this paleobiogeographic zone.

In Japan, the genus *Monodiexodina* has been reported with illustration of specimen from the lower part of the Kanokura Formation in the South Kitakami Belt (Fujimoto, 1956; Choi, 1973; Tazawa, 1976), the Kozaki Formation in the Kurosegawa Belt (Kanmera, 1963), the Moribu Formation of the Hida Gaien Belt (Tazawa et al., 1993), and the Takagami Conglomerate of the Choshi Peninsula, central Japan (Maeda and Mitsuoka, 1961). Among the *Monodiexodina*-bearing strata in Japan, the Moribu Formation and the Kanokura Formation

bear similar brachiopod faunas composed of mixed Boreal-, Tethyan-, and bipolar-type genera to that from the Oguradani Formation (Tazawa and Matsumoto, 1998). It is noteworthy that the Hida Gaaien Belt and the South Kitakami Belt also show close similarity in basic stratigraphy and faunal and floral successions throughout the Paleozoic (Tazawa, 2000).

Judging from the facts that the *Monodiexodina* from the Moribu Formation of the Hida Gaaien Belt and the Kanokura Formation of the South Kitakami Belt can be considered as the Murgabian in age (Ueno and Tazawa, 2003) along with that the Oguradani species is somewhat similar to *M. matsubaishi* (Fujimoto), the present Oguradani *Monodiexodina* would also be referable to the Murgabian (=Wordian, middle Middle Permian). Available brachiopod dating of the Oguradani Formation (Tazawa and Matsumoto, 1998) is also consistent with this age assignment. Moreover, the present report of *Monodiexodina* from the Oguradani Formation results in further reinforcing the original geologic (both geotectonic and paleobiogeographic) connection of the Hida Gaaien Belt with the South Kitakami Belt, which has been insisted by Tazawa (1989, 1996a, 1996b, 2000).

#### Acknowledgments

We are grateful to Mr. Yukio Miyake of Miya-mura, Gifu Prefecture, and Mr. Yosuke Ibaraki of Omi Natural History Museum for their field assistance. We also thank Dr. Isao Niikawa of the Department of Geology, Niigata University, for critical reviewing of the manuscript.

#### References

- Choi, D. R., 1973, Permian fusulinids from the Setamai-Yahagi district, Southern Kitakami Mountains, N. E. Japan. *Jour. Fac. Sci., Hokkaido Univ., Ser. 4*, **16**, 1-132.
- Fujimoto, H., 1956, A new species of *Parafusulina* from the Kitakami Massif, Japan. *Trans. Proc. Palaeont. Soc. Japan, N. S.*, no. 21, 157-160.
- Fujimoto, H., Kanuma, M. and Igo, H., 1962, On the Upper Paleozoic of the Hida Mountainland. In Fujimoto, H., ed., *Geologic Studies of the Hida Mountainland*, Shimada Seihan Insatsu, Tokyo, 44-70. (in Japanese)
- Hayasaka, I. and Matsuo, H., 1951, On the Permian fossils from Oguradani, Kamianama-mura, Ono-gun, Fukui Prefecture. *Jour. Geol. Soc. Japan*, **57**, 266. (in Japanese)
- Hayasaka, I. and Ozaki, K., 1955, A *Foordiceras* from the Permian formation of Central Japan. *Sci. Rep., Kanazawa Univ.*, **3**, 183-186.
- Horikoshi, E., Tazawa, J., Naito, N. and Kaneda, J., 1987, Permian brachiopods from Moribu, north of Takayama City, Hida Mountains, central Japan. *Jour. Geol. Soc. Japan*, **93**, 141-143. (in Japanese)
- Igo, H., 1957, Fusulinids of Fukuji, southern part of the Hida Massif, Central Japan. *Sci. Rep., Tokyo Kyoiku Daigaku, Sect. C*, **5**, 153-246.
- Igo, H., 1990, Paleozoic strata in the Hida "Gaaien" Belt. In Ichikawa, K., Mizutani, S., Hara, I., Hada, S. and Yao, A., eds., *Pre-Jurassic Terranes of Japan*, Nippon Insatsu Shuppan, Osaka, 41-48.
- Kanmera, K., 1963, Fusulines of the Middle Permian Kozaki Formation of southern Kyushu.



- Mem. Fac. Sci., Kyushu Univ., Ser. D*, **14**, 79-141.
- Kawai, M., Hirayama, K. and Yamada, N., 1957, *Explanatory Text of the Geologic Map of Japan, Scale 1:50,000 Arashimadake (Kanazawa, No. 58)*. Geological Survey of Japan, 110+13 p. (in Japanese)
- Maeda, S. and Mitsuoka, T., 1961, *Parafusulina matsubaishi* from the pebble of the Takagami Conglomerate in Choshi Peninsula, Chiba Prefecture. *Bull. Choshi Marine Lab., Chiba Univ.*, no. 3, 75-82.
- Niikawa, I., 1980, Geology and biostratigraphy of the Fukuji district, Gifu Prefecture, Central Japan. *Jour. Geol. Soc. Japan*, **86**, 25-36. (in Japanese)
- Niko, S., Yamakita, S., Otoh, S., Yanai, S. and Hamada, T., 1987, Permian radiolarians from the Mizuyagadani Formation in Fukuji area, Hida Marginal Belt and their significance. *Jour. Geol. Soc. Japan*, **93**, 431-433. (in Japanese)
- Ozaki, K., Yamada, K. and Kato, M., 1954, Permian System in Kamianama-mura, Ono-gun, Fukui Prefecture. *Jour. Geol. Soc. Japan*, **60**, 312-313. (in Japanese)
- Shi, G. R., Archbold, N. W. and Zhang, L. P., 1995, Distribution and characteristics of mixed (transitional) mid-Permian (Late Artinskian-Ufimian) marine faunas in Asia and their palaeogeographical implications. *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, **114**, 241-271.
- Tazawa, J., 1976, The Permian of Kesennuma, Kitakami Mountains: A preliminary report. *Earth Science (Chikyu Kagaku)*, **30**, 175-185.
- Tazawa, J., 1989, Comparison between the Palaeozoic formations of the Southern Kitakami Mountains, NE Japan and the Hida Marginal Belt, SW Japan: a preliminary note. *Earth Science (Chikyu Kagaku)*, **43**, 224-230. (in Japanese)
- Tazawa, J., 1991, Middle Permian brachiopod biogeography of Japan and adjacent regions in East Asia. In Ishii, K., Liu, X. Y., Ichikawa, K. and Huang, B. H., eds., *Pre-Jurassic Geology of Inner Mongolia, China: Report of China-Japan Cooperative Research Group, 1987-1989*, Matsuya Insatsu, Osaka, 213-230.
- Tazawa, J., 1996a, Permian continental shelf deposits in the Kitakami, Hida and Sikhote-Alin Mountains. *Chikyu Monthly (Gekkan Chikyu)*, **18**, 387-392. (in Japanese)
- Tazawa, J., 1996b, The Southern Kitakami and Hida Gaien belts in the Middle Permian in view of paleobiogeography. *Chikyu Monthly (Gekkan Chikyu)*, **18**, 730-734. (in Japanese)
- Tazawa, J., 2000, The Palaeozoic of the Hida Gaien, South Kitakami and Kurosegawa belts: correlation and tectonic history. *Mem. Geol. Soc. Japan*, **56**, 39-52. (in Japanese)
- Tazawa, J., 2001, Middle Permian brachiopods from the Moribu area, Hida Gaien Belt, central Japan. *Paleont. Res.*, **5**, 284-310.
- Tazawa, J. and Matsumoto, T., 1998, Middle Permian brachiopods from the Oguradani Formation, Ise district, Hida Gaien Belt, Central Japan. *Sci. Rep., Niigata Univ., Ser. E*, no. 13, 1-19.
- Tazawa, J., Tsusima, K. and Hasegawa, Y., 1993, Discovery of *Monodioxodina* from the Permian Moribu Formation in the Hida Gaien Belt, Central Japan. *Earth Science (Chikyu Kagaku)*, **47**, 345-348. (in Japanese)
- Ueno, K., 2003, The Permian fusulinoidean faunas of the Sibumasu and Baoshan blocks: their implications for the paleogeographic and paleoclimatologic reconstruction of the Cimmerian Continent. *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, **193**, 1-24.
- Ueno, K. and Tazawa, J., 2003, *Monodioxodina* from the Daheshen Formation, Jilin, Northeast China. *Sci. Rep., Niigata Univ., Ser. E*, no. 18, 1-16.
- Yamada, K., 1967, Stratigraphy and geologic structure of the Paleozoic formations in the upper Kuzuryu River district, Fukui Prefecture, Central Japan. *Sci. Rep., Kanazawa Univ.*, **12**, 185-207.
- Yamada, K., Ozaki, K., Kato, M., Yoshida, S. and Konishi, M., 1958, On Paleozoic formations in Izumi-mura, Ono-gun, Fukui Prefecture. *Jour. Geol. Soc. Japan*, **64**, 691. (in Japanese)
- Yamada, K. and Yamano, H., 1980, Find of Permian fossils from the Moribu Formation, Hida Mountains, Central Japan. *Sci. Rep., Kanazawa Univ.*, **25**, 53-65.