

## Lithostratigraphy of the Miocene Iwaine Formation in the Yatsuo area, central Japan

Raiki YAMADA\* and Naohiro YAMADA\*\*

### Abstract

We examined the Miocene Iwaine Formation formed in the margin of the Eurasian continent during the Japan Sea opening, of the Yatsuo Group in the Yatsuo area located in Toyama Prefecture, central Japan, on the basis of lithology and stratigraphy. The Iwaine Formation is composed mainly of monomictic autobreccia, polymictic volcanic breccia, pumice tuff and coherent andesites, with clastic rocks whose clasts were derived from the Iwaine Formation, in stratigraphically ascending order. Lithostratigraphy of the Iwaine Formation in the Yatsuo area will be of great significance to reveal the characteristics of volcanism during the Japan Sea opening.

*Key words:* Miocene, Iwaine Formation, Japan Sea opening, Yatsuo Group, lithostratigraphy.

### Introduction

The Japan Sea opening which is the formation of a back-arc basin in the margin of the Eurasian continent is the most significant event in the Neogene of Japan (e.g., Yoshida, 2009). Kano et al. (2007) suggested that the earliest stage of the Japan Sea opening had already initiated in latest Eocene. The Japan arc was then separated from the continent due to rapid spreading of the Japan Sea in 20–15 Ma (substage IA; Nakajima et al., 2015), and deposits during this stage are widely distributed in the Neogene strata (Fig. 1). Toyama Prefecture is

---


\* Graduate School of Science and Technology, Niigata University, Niigata 950-2181, Japan

\*\* Wakabayashi-green-haitsu #103, Gofuku-1-ku 5620, Toyama 930-0887, Japan

Corresponding author: R. Yamada,

e-mail: f17e069g@mail.cc.niigata-u.ac.jp

(Manuscript received 6 December, 2017; accepted 20 March, 2018)



**Fig. 1.** Map showing the distribution of Neogene sedimentary and volcanic rocks in Japan. Modified from Seamless digital geological map of Japan 1: 200000, Geological Survey of Japan.

one of the most suitable places to study the Japan Sea opening, because the strata formed during this period are well preserved without large deformation.

In Toyama Prefecture, there are two successions that were formed during the Japan Sea opening: the upper Oligocene stratum (Tori Conglomerate and Usunaka Moonstone Rhyolite) and the lower to middle Miocene Yatsuo Group (Fig. 2). The Tori Conglomerate and the Usunaka Moonstone Rhyolite were formed in the early stage of the opening on the margin of the continent. Since sedimentary environment of the lower to middle Miocene Yatsuo Group changed gradually from non-marine or shallow marine to deep sea (Hayakawa and Takemura, 1987), it is expected to reflect the deepening of the Japan Sea.

The Iwaine Formation of the Yatsuo Group and equivalent beds, widely distributed in the Hokuriku Region, central Japan (Fig. 3), are composed mainly of andesites which erupted in the margin of the Eurasian continent in Early Miocene (Takahashi and Shuto, 1999). Petrological studies of the Iwaine Formation have been conducted by Ishiwatari and




Fig. 2. Stratigraphy of strata associated with the Japan Sea opening in Toyama Prefecture (after Sudo (1979), and Hayakawa and Takemura (1987)). The youngest and oldest ages are shown as age data of each formation (1, Ueda and Aoki, 1970; 2, Shibata, 1973; 3, Ganzawa, 1983; 4, Nakajima and Mizushima, 1984; 5, Japan National Oil Corporation, 1985; 6, Hayakawa and Danhara, 1986). The tectonic interpretations associated with the Japan Sea opening are based on Hayakawa and Takemura (1987; 7), Fournier et al. (1994; 8), Ishida et al. (1998; 9), Takahashi and Shuto (1999; 10), Tamaki et al. (2006; 11), Ayalew and Ishiwatari (2011; 12), Sato et al. (2013; 13). K-Ar: K-Ar age. FT: zircon fission track age. Gp.: Group. Fm.: Formation.




Fig. 3. Index map showing the distribution of Neogene basalts and andesites in Hokuriku Region (modified from Seamless digital geological map of Japan 1: 200,000, Geological Survey of Japan) and the study area. Pref.: Prefecture.

Ohama (1997), Takahashi and Shuto (1999), Tsuchihashi and Ishiwatari (2006), Ishiwatari et al. (2007), Sato et al. (2013), and Watanabe and Takahashi (2017). Most petrological studies concluded that the Iwaine Formation was formed by volcanic activities associated with hot mantle wedge in the margin of the Eurasian continent, in a subduction zone setting. However, this state differed from the present subduction and volcanic activities in the Japan arc. Although many researchers have referred to lithology and stratigraphy of the Iwaine Formation (e.g., Tsuda, 1953, 1955; Sakamoto and Nozawa, 1960; Hayakawa and Takemura, 1987), detailed studies on the lithostratigraphy of the Iwaine Formation are still not enough. Geology, including lithology and stratigraphy, should be as significant as petrology to reveal the characteristics of volcanism of the Iwaine Formation. Therefore, we examined lithology and stratigraphy of the Iwaine Formation in the Yatsuo area (Fig. 3), and accordingly we report the results of our field research and figure out some unsolved issues on stratigraphy of the Iwaine Formation.

### **Regional geology in the Yatsuo area**

The stratigraphic division of the Neogene in the Yatsuo area was first founded by Makiyama (1930), and many other researchers have revised it (e.g., Imamura, 1936, 1937; Fujita and Nakagawa, 1948; Tsuda and Chiji, 1950; Tsuda, 1953; Nakaseko, 1953, 1954; Sakamoto et al., 1959; Sakamoto and Nozawa, 1960; Hayakawa and Takemura, 1987). Hayakawa and Takemura (1987) showed a stratigraphic division based on the correlation of key tuff beds and detailed columnar sections of the Miocene in the Yatsuo area. We think that their stratigraphic division is reasonable because they associated lithostratigraphy of Neogene strata in the Yatsuo area with the Japan Sea opening, and hence follow it in our study. However, type localities of some formations have already sunk into rivers due to construction of dams.

Mesozoic basement rocks are distributed in the southern part of the target area, and the Miocene Yatsuo Group overlies them (Fig. 4); the Miocene Yatsuo Group is composed of the Nirehara, Iwaine, Iozen, Kurosedani and Higashibessho formations, in stratigraphically ascending order (Fig. 5). The Yatsuo Group strikes ENE–WSW and dips 10–40° N (Fig. 4).

#### **1. Basement rocks**

The basement rocks of the Miocene in the Yatsuo area consist of the Hida Granitic Rocks (Kano, 1990), the Tetori and Jinzu groups (Matsukawa et al., 2014). The granite in this area is weathered and appears to be pale pink, consisting of potassium feldspar, plagioclase, quartz, biotite, and opaque minerals. The Tetori and Jinzu groups are in fault contact with the Hida Granitic Rocks (e.g., Sakamoto and Nozawa, 1960). The Tetori Group in the study area consists of upper Oxfordian marine sandstone and mudstone, whereas the Jinzu Group consists of Lower Cretaceous non-marine sandstone and mudstone (Matsukawa et al., 2014).




Fig. 4. Geological map and two cross sections of the Yatsuo area. Fm.: Formation.

## 2. Nirehara Formation

**Name** Named by Tsuda and Chiji (1950).

**Type locality** The bank of the Jinzugawa River in Nirehara, Toyama City, Toyama Prefecture (Hayakawa and Takemura, 1987).







Age	Formations	Column	Rock type
Early to middle Miocene	Higashibessho Fm.		Massive mudstone
	Kurosedani Fm.		Conglomerate, sandstone and mudstone
	Iozen Fm.		Rhyolitic pyroclastic rocks
	Iwaine Fm.		Andesitic volcanic and volcanoclastic rocks
	Nirehara Fm.		Sandstone and conglomerate
Meso-zoic	Tetori and Jinzu Gps. Hida Granitic Rocks		Sandstone, mudstone and granite

Fig. 5. Stratigraphy of the study area. The division of basement rocks is after Kano (1990) and Matsukawa et al. (2014). Gp.: Group. Fm.: Formation.

**Distribution** The Nirehara Formation is distributed narrowly between Taniori and Kiridani, southern part of the study area (Fig. 4).

**Thickness** 100–250 m.

**Stratigraphic relationship** The Nirehara Formation is in unconformable or fault contact with the basement rocks in the study area (Fig. 4).

**Lithology** The lower part of the Nirehara Formation is composed mainly of matrix-supported conglomerate containing rounded pebbles of granites and chert (approximately 10 cm in diameter). The conglomerate rarely shows imbricate structure and includes lenticular sandstone and mudstone. The upper part is composed of massive (rarely showing cross bedding) arkose sandstone and mudstone.

**Fossil** Tsuda (1953), Sakamoto and Nozawa (1960), and Hayakawa and Takemura (1987) reported unidentifiable fragments of plant fossils. Kashiwagi (2012) reported radiolarian fossils from chert gravels in the formation and concluded that the chert gravels were originated from the chert in the Tamba–Mino Belt or the chert gravels in the conglomerate beds of the Tetori Group.

### 3. Iwaine Formation

**Name** Named by Imamura (1937).

**Type locality** The bank of the Jinzu River in Iwaine, Toyama City, Toyama Prefecture (Hayakawa and Takemura, 1987).

**Distribution** The Iwaine Formation is distributed widely between Nunotani and Mt. Otakayama, the southern part of the study area (Fig. 4).

**Thickness** 600–1450 m.

**Stratigraphic relationship** The Iwaine Formation conformably covers the Nirehara Formation (Fig. 5).

**Lithology** The Iwaine Formation consists of coherent andesites, volcanoclastic and clastic rocks. Volcanoclastic and clastic rocks are predominant in the lower part of the formation, whereas coherent andesites are predominant in the upper part. Coherent andesites are generally massive but partly has joints. Thin beds of volcanoclastic or clastic rocks are intercalated often between coherent andesites. Volcanoclastic rocks in the Iwaine Formation are composed mainly of autobreccia, volcanic breccia and pumice tuff showing pale to dark green color.

**Fossil** Tsuda (1953) reported unidentifiable fragments of plant fossils. Sakamoto and Nozawa (1960) reported molluscan (e.g., *Soletollina* sp., *Calyptrea* sp. and *Nassarius* sp.) and plant (*Fagus antipofii*) fossils from mudstone in the formation. In this study, a gastropod molluscan shell of *Cerithideopsilla* sp. was found from siltstone in the upper part of the formation (Fig. 7).

**Age** K–Ar and FT (fission track) ages have been reported by Shibata (1973; K–Ar age;  $13.6 \pm 0.7$ ,  $15.9 \pm 0.9$ ,  $16.0 \pm 0.9$  Ma), Japan National Oil Corporation (1985; K–Ar age;  $15.8 \pm 1.1$ ,  $18.9 \pm 0.9$  Ma), Kaneko (2001; K–Ar age;  $16.3 \pm 0.9$ ,  $16.5 \pm 0.9$  Ma), and Itoh and Watanabe (2006; FT age;  $17.2 \pm 0.9$  Ma). Recently, Nakajima et al. (2015) reported the weighted average of  $^{238}\text{U}$ – $^{206}\text{Pb}$  ages of  $17.6 \pm 0.3$  Ma from zircons in the welded tuff of the Iwaine Formation in the Kubusugawa basin.

#### 4. Iozen Formation

**Name** Named by Fujita and Nakagawa (1948).

**Type locality** Mt. Iozen on the boundary between Nanto City, Toyama Prefecture and Kanazawa City, Ishikawa Prefecture (Hayakawa and Takemura, 1987).

**Distribution** The Iozen Formation is distributed narrowly around Yomedani, the westernmost part of the study area (Fig. 4).

**Thickness** Less than 50 m in this area.

**Stratigraphic relationship** The Iozen Formation conformably covers the Iwaine Formation (Fig. 5).

**Lithology** The Iozen Formation in the study area is composed of rhyolitic pumice tuff and lapilli tuff. The Iozen Formation can be distinguished from the Iwaine Formation by presence of rhyolitic rocks.

**Age** K–Ar and FT ages have been reported by Shibata (1973; K–Ar age;  $14.1 \pm 1.2$  Ma), Ganzawa (1983; FT age;  $15.0 \pm 2.1$ ,  $16.5 \pm 1.4$  Ma) and Kaneko (2001; K–Ar age;  $14.3 \pm 0.3$ ,  $14.4 \pm 0.3$  Ma).

#### 5. Kurosedani Formation

**Name** Named by Tsuda and Chiji (1950).

**Type locality** Area between Kakehata and Do, Toyama City, Toyama Prefecture (Hayakawa and Takemura, 1987).

**Distribution** The Kurosedani Formation is distributed widely between Do and Yomedani, the northern part of the study area (Fig. 4).

**Thickness** More than 900 m.

**Stratigraphic relationship** The Kurosedani Formation unconformably covers the Iozen and Iwaine formations (Fig. 5).

**Lithology** The lower part of the formation consists mainly of irregularly alternating beds of conglomerate, sandstone, and mudstone, whereas the upper part consists mainly of mudstone and alternating beds of sandstone and mudstone. Conglomerate contains rounded to sub-rounded pebbles of andesite. The lithofacies of the Kurosedani Formation changes gradually into the Higashibessho Formation.

**Fossil** The Kurosedani Formation is a Miocene fossiliferous formation in Japan. Various bivalve and gastropod fossils have been found including *Anadara kakehataensis*, *Cerithideopsilla* n. sp., *Vicarya yokoyamai*, and *Vicaryella ishiiana* var. (Tsuda, 1953, 1959). Tsuda (1953) also reported fossils of plants, Echinodermata, Crustacea, coral and shark teeth from the formation.

**Age** K-Ar and FT ages have been reported by Hayakawa (1983; FT age;  $16.0 \pm 2.4$  Ma), Nakajima and Mizushima (1984; FT age;  $15.0 \pm 1.0$  Ma), Hayakawa and Danhara (1986; FT age;  $16.8 \pm 0.9$  Ma), Kaneko (2001; K-Ar age;  $13.8 \pm 0.4$ ,  $14.0 \pm 0.4$  Ma) from tuffs of the Kurosedani Formation. Yamada et al. (1998), Kaneko (2001), and Itoh and Watanabe (2006) reported 15.3–12.2 Ma of K-Ar and FT ages from volcanic rocks in the Fukuhira Formation, which is heterotopic with the Kurosedani Formation in the eastern part of Toyama Prefecture.

## 6. Higashibessho Formation

**Name** Named by Fujita and Nakagawa (1948).

**Type locality** Higashibessho, Tonami City, Toyama Prefecture (Hayakawa and Takemura, 1987).

**Distribution** The Higashibessho Formation is distributed around Kashio, the northern part of the study area (Fig. 4).

**Thickness** More than 500 m.

**Stratigraphic relationship** The Higashibessho Formation conformably covers the Kurosedani Formation in the study area (Fig. 5) and is unconformably covered by the Upper Miocene Otogawa Formation, the Tonami Group outside the study area (Hayakawa and Takemura, 1987).

**Lithology** The Higashibessho Formation is composed mainly of massive mudstone showing dark gray color. Thin conglomerate beds (2–10 cm thick) are intercalated in the formation and often include allochthonous bivalve and gastropod fossils.

**Fossil** The Higashibessho Formation abundantly yields fossils of mollusk, foraminifer, radiolarian and diatom (Hayakawa and Takemura, 1987).



### Lithostratigraphy of the Iwaine Formation

We obtained columnar sections along fifteen routes in the Iwaine Formation (Fig. 6). We classify volcanoclastic rocks after the definition of Fisher (1961, 1966). Following the lithology and stratigraphy, the distribution of lithofacies can be classified into four parts (Fig. 7).

Monomictic variegated autobreccia (Fig. 8a), whose clasts are composed completely of amphibole-bearing andesite (Fig. 8b), without matrix occupies the lowest part of the Iwaine Formation (Fig. 7), and its maximum thickness reaches approximately 250 m. Coherent amphibole-bearing andesite sometimes occurs with the autobreccia.

Matrix supported volcanic breccia (Fig. 8c and d) composed of poorly sorted polymictic andesitic clasts overlie the autobreccia in many routes (Fig. 7). Generally, the volcanic breccia shows massive or reverse-grading structures (Fig. 8d). Its thickness indicates greatly lateral change. Clasts in the volcanic breccia consist of sub-angular to sub-rounded gravels (mainly 30–50 cm; maximum 100 cm in diameter) of various andesites. Conglomerate, sandstone and mudstone whose andesitic clasts derived from the Iwaine Formation overlie thickly the volcanic breccia in Routes 7 and 13 (Fig. 8e).

In some routes, poorly sorted massive pumice tuff (Fig. 8f) overlies the clastic rocks or the polymictic volcanic breccia (Fig. 7). The pumice tuff is composed of lithic clasts of andesites and pumice (approximately 5 cm in diameter), with tuffaceous matrix. The pumice




Fig. 6. Map showing place-names and routes along which columnar sections were obtained. Alphabets (a-p) with arrows indicate sampling points and locality of outcrops in Fig. 8. Riv.: River.




Fig. 7. Correlated columnar sections of the Iwaine Formation, the Yatsuo area and simplified distribution of rock types. Routes are shown in Fig. 6. Fm.: Formation.


tuff is partly welded in Route 13 (Fig. 8g and h). The content of lithic clasts and pumice changes with stratigraphic horizons and routes. In Route 8, the pumice tuff includes a lot of lithic clasts instead of pumice, while the pumice tuff in other routes includes much pumice rather than lithic clasts.

Coherent andesites (Fig. 8i–l) overlie thickly the pumice tuff or breccia (autobreccia and volcanic breccia) of the lower part. Pyroxene phenocrysts (two pyroxene, clinopyroxene or orthopyroxene; Fig. 8j) occur in coherent andesites of the upper part of the Iwaine Formation. Coherent andesites imply often columnar or platy joints (Fig. 8k). Coherent andesites change gradually to homogeneous autobreccia upward, which comprises reddish-brown colored angular blocks or vesicular scoriaceous clasts (Fig. 8l). Clastic rocks and polymictic volcanic breccia showing various lithofacies are intercalated usually between coherent andesites. A bed (10 m in thick) of very well sorted monomictic lapillistone occur in Route 5 (Fig. 8m). A gastropod molluscan shell of *Cerithideopsis* sp. (Fig. 8n) was found from clastic rocks (Fig. 8o) of the uppermost Iwaine Formation, Route 13 in this study. The locality is probably the same as the locality where Sakamoto and Nozawa (1960) found some fossils including *Cerithidea miofluviatilis*. An outcrop, where a dike intrudes in volcanic breccia and partly shows red color, is exposed in Route 4 (Fig. 8p). The dike changes to effusive coherent andesite in the outcrop. Massive aphyric andesite is distributed successively in the top of the Iwaine Formation.


### Implication

The Yatsuo Group is associated with the Japan Sea opening (Hayakawa and Takemura, 1987; Kaneko, 2001). Therefore, many experts have studied stratigraphy especially in the Yatsuo area, the central part of Toyama Prefecture, as already mentioned in the introduction. Following the stratigraphy in the Yatsuo area, some ones have studied in the eastern part (e.g., Nozawa and Sakamoto, 1960; Sumi and Nozawa, 1973; Kaneko, 2001) and the western part (e.g., Inoue et al., 1964; Sudo, 1979; Ganzawa, 1983) of Toyama Prefecture. Unfortunately, some formations which are not observed in the Yatsuo area occur in these areas (e.g., Fukuhira Formation in the eastern part; Tori Conglomerate and Usunaka Moonstone Rhyolite in the western part). In addition, the Neogene strata in some areas have been classified according to the division of the Hokuriku Group (Sakamoto et al., 1959), which had been widely accepted by geologists studying the Neogene in the Hokuriku Region until Hayakawa and Takemura (1987) proposed the division of the Yatsuo Group. As a result, the division and stratigraphy of the Yatsuo Group is waiting to be solved.

Sumi and Nozawa (1973) classified the Iwaine Formation in the Uozu area into three members. Although Kaneko (2001) correlated the Iwaine Formation in the Uozu area to the upper Iozen Formation, it includes both of andesites and rhyolites (Kaneko, 2001). Since the



**Fig. 8.** Photographs of outcrops, a float, a hand specimen and thin sections (Pl: plagioclase, Opx: orthopyroxene, V: vesicle, Amp: amphibole). Localities of outcrops and sampling points are shown in Fig. 6. **a**, a float of monomictic autobreccia. **b**, a thin section of monomictic autobreccia. Broken lines indicate boundaries, which are unclear, of clasts. **c**, a cross section of polymictic volcanic breccia showing massive structure. Hammer for scale. **d**, a cross section of reverse-graded polymictic volcanic breccia. Hammer for scale. **e**, a cross section of clastic rocks whose all clasts are derived from the Iwaine Formation. **f**, a cross section of massive pumice tuff. **g**, a cross section of the contact between welded and non-welded parts of pumice tuff. **h**, a hand specimen of welded tuff. **i**, a cross section of massive coherent andesite. **j**, a thin



section of orthopyroxene-bearing andesite. **k**, a cross section of aphyric andesite showing columnar and platy joints. Parson for scale. **l**, a cross section of polymictic autobreccia consisting of andesitic coherent blocks. It would be clinker (flow breccia). Hammer (circle) for scale. **m**, a cross section of very well-sorted monomictic lapillistone. **n**, a photograph of a gastropod mollusk of *Cerithideopsis* sp. obtained from clastic rocks (o), the uppermost Iwaine Formation. **o**, a cross section of fossiliferous clastic rocks, where a gastropod mollusk of *Cerithideopsis* sp. was found (n). **p**, a cross section of a dike intruding host deposits (volcaniclastic rocks). The host deposits adjacent to the dike show red color by high-temperature oxidation.

Iwaine and Iozen formations comprise of andesites and rhyolites (e.g., Hayakawa and Takemura, 1987) respectively, it is difficult to judge which is correct. Nevertheless, Ganzawa (1983) and Takahashi and Shuto (1999) reported that amphibole-bearing andesites occur in the lowest part of the Iwaine Formation and that pyroxene-bearing andesites overlie them. The lithostratigraphy of the Iwaine Formation in the study area is consistent with their reports.

The stratigraphic division of volcanic rocks in the Yatsuo Group is very important to reveal the temporal change of characteristics of volcanism during the Japan Sea opening. Accordingly, lithostratigraphic evidence reported by this study will be significant to solve such issues. What we need to do is establishing formative history of the Iwaine Formation based on facies analysis, which is necessary to propose the stratigraphic division of this formation.

### Acknowledgments

We especially thank Mr. Mitsuhiro Nagata of the University of Toyama for his advice to write this paper and for accompanying us several times in the field. Helpful comments by two reviewers (Associate Professor Hayato Ueda and Professor Norie Fujibayashi of Niigata University) and Professor Atsushi Matsuoka of Niigata University greatly improved the manuscript. Proofreading of the English manuscript by Professor M. Satish Kumar of Niigata University and Dr. Yiyi Zhang of China University of Geosciences (Beijing) is greatly appreciated. Our sincere gratitude goes to Associate Professor Toshiro Takahashi and Associate Professor Kyoko Kataoka of Niigata University for discussing this study and for sharing their knowledge on volcanic rocks. The authors are also grateful to Professor Shigeru Otoh of the University of Toyama for his pointing out imperfection in logic and sharing his knowledge on Miocene strata in Japan. We thank Dr. Kazuo Kaneko for sharing his knowledge of fossils from the Iwaine Formation with us, and Mr. Taiji Hara for allowing us to stay in the temple for the parishioners during our geological survey.

### References

- Ayalew, D. and Ishiwatari, A., 2011, Comparison of rhyolites from continental rift, continental arc and oceanic island arc: Implication for the mechanism of silicic magma generation. *Island Arc*, **20**, 78–93.
- Fisher, R. V., 1961, Proposed classification of volcanoclastic sediments and rocks. *Geol. Soc. Amer. Bull.*, **72**, 1409–1414.
- Fisher, R. V., 1966, Rocks composed of volcanic fragments and their classification. *Earth-Sci. Rev.*, **1**, 287–298.
- Fournier, M., Jolivet, L. and Huchon, P. (1994). Neogene strike-slip faulting in Sakhalin and the Japan Sea opening. *Jour. Geophys. Res.*, **99**, 2701–2725.
- Fujita, K. and Nakagawa, C., 1948, The Tertiary in the Tonami area, Toyama Prefecture. *54th Ann. Meet. Geol. Soc. Japan, Abstr.*, 125 (in Japanese\*).
- Ganzawa, Y., 1983, "Green Tuff" Movement defined by fission track ages of igneous rocks Part 2: Futomiyama area of Toyama Prefecture, central Japan. *Jour. Geol. Soc. Japan*, **89**, 271–286 (in Japanese with English abstract).



- Geological Survey of Japan (GSJ). Seamless digital geological map of Japan 1:200,000. May 29, 2015 version. <https://gbank.gsj.jp/seamless/seamless2015/2d/>.
- Hayakawa, H., 1983, Stratigraphy and geochronology of the Neogene in the west part of the Yatsuo are, Toyama Prefecture, Central Japan. *News Osaka Micropaleontol.*, no. 10, 1-13 (in Japanese with English abstract).
- Hayakawa, H. and Danhara, T., 1986, Fission-track Age of a Tuff Layer intercalated in the Kurosedani Formation, Yatuso Area, Central Japan. *News Osaka Micropaleontol.*, no. 14, 63-69 (in Japanese with English abstract).
- Hayakawa, H. and Takemura, A., 1987, The Neogene System in the Yatsuo area, Toyama Prefecture, central Japan. *Jour. Geol. Soc. Japan*, **93**, 717-732 (in Japanese with English abstract).
- Imamura, S., 1936, Outline of geology of Toyama Prefecture. *Toyama Hakubutsu Gakkaishi* (Journal of Toyama Natural Historic Society; English translation from the original written in Japanese), no. 1, 1-25 (in Japanese\*).
- Imamura, S., 1937, Geology of the Jinzugawa basin. *Tomiko Hakubutsu Dokokaishi* (Journal of Natural Historic Association of Toyama High School; English translation from the original written in Japanese), no. 3, 1-11 (in Japanese\*).
- Inoue, M., Mizuno, A. and Nozawa, T., 1964, *Johana, Explanatory text of geological map of Japan, Scale 1:50,000*. Geological Survey of Japan, AIST, Tsukuba, 32 p. (in Japanese with English abstract).
- Ishida, H., Ishiwatari, A. and Kagami, H., 1998, The Mt. Wasso moonstone rhyolitic welded tuff in the Neogene Hokuriku Group, central Japan. *Jour. Geol. Soc. Japan*, **104**, 281-295 (in Japanese with English abstract).
- Ishiwatari, A. and Ohama, H., 1997, Clinopyroxene basalt dikes in the Miocene Iwaine Formation, Hokuriku Province, Japan: Various continental arc magmas including shoshonite series and origin of the clinopyroxene phenocrysts. *Jour. Geol. Soc. Japan*, **103**, 565-578 (in Japanese with English abstract).
- Ishiwatari, A., Tsuchihashi, H., Shuto, K. and Sato, M., 2007, Sr-isotopic ratio of the Miocene adakite in western Toyama Prefecture, central Japan and Sr content of its plagioclase. *114th Ann. Meet. Geol. Soc. Japan, Abstr.*, 357 (in Japanese).
- Itoh, Y. and Watanabe, M., 2006, Fission-track dating of the Lower to Middle Miocene in the Uozu area, central Japan. *Bull. Geol. Surv. Japan*, **57**, 57-59 (in Japanese with English abstract).
- Japan National Oil Corporation, 1985, Survey report of MITI "Toyama" well, 60 p. (in Japanese\*).
- Kaneko, K., 2001, Stratigraphy and geotectonic history of Miocene volcanic-volcaniclastic rocks in the eastern part of Toyama Prefecture, central Japan. *Jour. Geol. Soc. Japan*, **107**, 729-748 (in Japanese with English abstract).
- Kano, K., Uto, K. and Ohguchi, T., 2007, Stratigraphic review of Eocene to Oligocene successions along the eastern Japan Sea: Implication for early opening of the Japan Sea. *Jour. Asian Earth Sci.*, **30**, 20-32.
- Kano, T., 1990, Intrusive relation of the Okumayama Granitic Mass (Shimonomoto type) into the Iori Granitic Mass (Funatsu type) in the Hayatsukigawa area; Re-examination of the sub-division for early Mesozoic granites (Funatsu Granites) in the Hida region. *Jour. Geol. Soc. Japan*, **96**, 379-388 (in Japanese with English abstract).
- Kashiwagi, K., 2012, Sedimentary environment of the Nirehara Formation (Lower Miocene) in the Yatsuo area, Toyama Prefecture of central Japan and Paleozoic and Mesozoic radiolarian fossils from chert pebbles and cobbles. *Mem. Fukui Pref. Dinosaur Mus.*, **11**, 27-47 (in Japanese with English abstract).
- Makiyama, J., 1930, Outline of the Neogene in Ishikawa and Toyama prefectures. *Chikyū*, **14**, 161-174 (in Japanese\*).
- Matsukawa, M., Fukui, M., Ogawa, Y., Tago, T., Koarai, K., Ohira, H. and Hayashi, K., 2014, Stratigraphy of the Tetori Group and Jinzu Group (new name) in Gifu and Toyama prefectures, central Japan. *Jour. Geol. Soc. Japan*, **120**, 147-164.
- Nakajima, T., Danhara, T., Iwano, H. and Yamashita, T., 2015, Opening of the Sea of Japan and tectonics of Japanese islands during Green Tuff Epoch. *122nd Ann. Meet. Geol. Soc. Japan, Abstr.*, 23 (in Japanese).
- Nakajima, T. and Mizushima, S., 1984, Fission-track ages of *Miogypsina-Operculina* horizons in Hokuriku District, central Japan. *Jour. Geol. Soc. Japan*, **90**, 667-670 (in Japanese).
- Nakaseko, K., 1953, On the Stratigraphy of the Upper Neogene Formation in the Southern Part of Toyama Prefecture. *Osaka Daigaku Nankou Osaka Daigaku Hokkou Rika Houkoku* (Science Report of South

- Collage and North Collage of Osaka University; English translation from the original written in Japanese), **2**, 87–100 (in Japanese with English abstract\*).
- Nakaseko, K., 1954, Correction of the Stratigraphy of the Upper Neogene Formation in The Southern Toyama Prefecture. *Osaka Daigaku Nankou Osaka Daigaku Hokkou Rika Houkoku* (Science Report of South Collage and North Collage of Osaka University; English translation from the original written in Japanese), **3**, 97–105 (in Japanese with English abstract\*).
- Nozawa, T. and Sakamoto, T., 1960, *Gohyakkoku, Explanatory text of geological map of Japan, Scale 1:50,000*. Geological Survey of Japan, AIST, Tsukuba, 68 p. (in Japanese with English abstract).
- Sakamoto, T., Imai, I., Mizuno, A., Sumi, Y. and Inoue, M., 1959, Cenozoic Strata in the Southern Part of the Toyama Sedimentary Basin. *Bull. Geol. Surv. Japan*, **10**, 75–82 (in Japanese with English abstract).
- Sakamoto, T. and Nozawa, T., 1960, *Yatsuo, Explanatory text of geological map of Japan, Scale 1:50,000*. Geological Survey of Japan, AIST, Tsukuba, 69 p. (in Japanese with English abstract).
- Sato, M., Shuto, K., Uematsu, M., Takahashi, T., Ayabe, M., Takanashi, K., Ishimoto, H. and Kawabata, H., 2013, Origin of late Oligocene to middle Miocene Adakitic andesites, high magnesian andesites and basalts from the Back-arc margin of the SW and NE Japan arcs. *Jour. Petrol.*, **54**, 481–524.
- Shibata, K., 1973, K–Ar ages of volcanic rocks from the Hokuriku Group. *Mem. Geol. Soc. Japan*, no. 8, 143–149 (in Japanese with English abstract).
- Sudo, S., 1979, On the geology of the Futomiyama Mountains, Toyama Prefecture, central Japan (Preliminary report). *Mem. Geol. Soc. Japan*, no. 17, 187–194 (in Japanese with English abstract).
- Sumi, Y. and Nozawa, T., 1973, *Uozu, Explanatory text of geological map of Japan, Scale 1:50,000*. Geological Survey of Japan, AIST, Tsukuba, 104 p. (in Japanese with English abstract).
- Takahashi, T. and Shuto, K., 1999, Genesis of adakitic andesite, high-magnesian andesite, calc-alkaline andesite and tholeiitic andesite in the Miocene Iwaine Formation, southern part of Toyama Prefecture, Japan. *Jour. Geol. Soc. Japan*, **105**, 789–809 (in Japanese with English abstract).
- Tamaki, M., Itoh, Y. and Watanabe, M., 2006, Paleomagnetism of the lower to middle Miocene series in the Yatsuo area, eastern part of southwest Japan: Clockwise rotation and marine transgression during a short period. *Bull. Geol. Surv. Japan*, **57**, 73–88.
- Tsuchihashi, H. and Ishiwatari, A., 2006, Adakitic hornblende-andesite and xenoliths in the Miocene volcanic rocks, western part of Toyama Prefecture, central Japan. *113rd Ann. Meet. Geol. Soc. Japan, Abstr.*, 243 (in Japanese).
- Tsuda, K., 1953, Geology of Yatsuo Area in Toyama Prefecture. *Jour. Fac. Sci. Niigata Univ., Ser. II*, **1**, 1–35 (in Japanese with English abstract).
- Tsuda, K., 1955, On the sedimentary environment of the Yatsuo Group: A study of the sedimentary environment of the Miocene deposits in the so-called “Green Tuff” Region (Part 1). *Jour. Geol. Soc. Japan*, **61**, 532–542 (in Japanese with English abstract).
- Tsuda, K., 1959, New Miocene molluscs from the Kurosedani Formation in Toyama Prefecture, Japan. *Jour. Fac. Sci. Niigata Univ., Ser. II*, **3**, 67–137.
- Tsuda, K. and Chiji, M., 1950, Geological problems in the Yatsuo area. *56th Ann. Meet. Geol. Soc. Japan, Abstr.*, 304 (in Japanese).
- Ueda, Y. and Aoki, K., 1970, K–Ar dating on moonstone rhyolite in Toyama Prefecture, central Honshu. *Jour. Japanese Assoc. Mineral. Petrol. Econ. Geol.*, **63**, 28–29 (in Japanese with English abstract).
- Watanabe, S. and Takahashi, T., 2017, Geological and petrological study of Miocene volcanic rock in Tateyama Kamiichi area eastern part of Toyama Prefecture. *124th Ann. Meet. Geol. Soc. Japan, Abstr.*, 180 (in Japanese).
- Yamada, N., Sakamoto, T. and Kaneko, K., 1998, K–Ar ages of “the Takamineyama Volcanic Rocks”, Toyama Prefecture, central Japan. *Earth Sci. (Chikyu Kagaku)*, **52**, 235–239 (in Japanese).
- Yoshida, T., 2009, Late Cenozoic magmatism in the northeast Honshu arc, Japan. *Earth Sci. (Chikyu Kagaku)*, **63**, 269–288 (in Japanese with English abstract).

\*The title was translated by the authors.