

Updated radiolarian zonation for the Jurassic in Japan and the western Pacific

Atsushi MATSUOKA* and Tsuyoshi ITO**

Abstract

An updated version from radiolarian zonal scheme in the 1990s for the Jurassic in Japan and the western Pacific is presented and vertical distributions of selected radiolarian species are illustrated. The major modifications are as follows: The *Bipedis horiae* Zone (JR0) is added below the *Parahsuum simplum* Zone (JR1). The base of the *Bipedis horiae* Zone (JR0) is defined by the last occurrence biohorizon of *Haeckelicyrtium breviora* Sugiyama and corresponds to the Triassic–Jurassic boundary. The base of the *Parahsuum simplum* Zone is dated within the Sinemurian. The *Tricolocapsa plicarum* Zone (JR4), *Tricolocapsa conexa* Zone (JR5), *Stylocapsa(?) spiralis* Zone (JR6), and *Pseudodictyomitra primitiva* Zone (JR8) are modified to the *Striatojaponocapsa plicarum* Zone (JR4), *Striatojaponocapsa conexa* Zone (JR5), *Kilinora spiralis* Zone (JR6), and *Loopus primitivus* Zone (JR8), respectively, in accordance to the change of generic assignment of zone-nominal species.

Key words: Jurassic, radiolaria, zonation, revision, Japan, western Pacific.

Introduction

Radiolarian zonation for the Jurassic and Lower Cretaceous using data from land sections in Japan and deep sea cores in the western Pacific was proposed in Matsuoka (1995a). Since then, taxonomic and biostratigraphic studies on Jurassic radiolarians have been actively carried out in Europe (e.g., Steiger, 1992; O'Dogherty et al., 2006, 2017), North

* Department of Geology, Faculty of Science, Niigata University, Niigata 950-2181, Japan

** Research Institute of Geology and Geoinformation, Geological Survey of Japan, AIST, Tsukuba 305-8567, Japan
Corresponding author: A. Matsuoka,

amatsuoka@geo.sc.niigata-u.ac.jp

(Manuscript received 20 December, 2018; accepted 15 March, 2019)

America (e.g., Hull, 1997; Carter et al., 1998; Yeh, 2009; Pessagno et al., 2009), Asia (e.g., Sugiyama, 1997; Hori, 1999; Matsuoka et al., 2002), and Antarctica (Kießling, 1999). International joint research on taxonomy and biostratigraphy of Jurassic radiolarians has been performed continuously (Carter et al., 2010; Gorican et al., 2006, 2013, 2018; O'Dogherty et al., 2009). The accumulation of new data in the last two decades requires a revision of the zonation and age assignments. Although revision of the zonation was briefly reported in Matsuoka (2006), detailed explanations have not been demonstrated yet. An updated version of the zonal scheme of Matsuoka (1995a) for the entire Jurassic in Japan and the western Pacific is presented in this paper.

A brief review of Jurassic radiolarian zonation in Japan and the western Pacific

Mesozoic radiolarian studies in Japan were carried out to clarify the geology of accretionary complexes since the 1970s. In the early 1980s several research groups started radiolarian biostratigraphic work at almost the same time so that several independent zonal schemes were proposed. A historical review on Jurassic radiolarian biostratigraphic research in Japan is given in Gorican et al. (2018).

Interval zones for the entire Jurassic were established by Matsuoka and Yao (1986). Deep sea cores in the western Pacific obtained from Leg 129 of the Ocean Drilling Program (ODP) contributed to testifying the zonal scheme for the Middle and Upper Jurassic established in Japan (Matsuoka, 1992). Combining data from land sections in Japan with data from the ODP materials, Matsuoka (1995a) proposed a zonal scheme for the Jurassic and Lower Cretaceous. A detailed data set for this zonation was presented in Matsuoka (1995b). The Jurassic succession includes the *Parahsuum simplum* Zone (JR1), *Trillus elkhornensis* Zone (JR2), *Laxtorum(?) jurassicum* Zone (JR3), *Tricolocapsa plicarum* Zone (JR4), *Tricolocapsa conexa* Zone (JR5), *Stylocapsa(?) spiralis* Zone (JR6), *Hsuum maxwelli* Zone (JR7), and *Pseudodictyomitra primitiva* Zone (JR8) in ascending order. In addition to these zones, the *Pseudodictyomitra carpatica* Zone (KR 1) includes the uppermost Jurassic as well. Most of these zones are introduced in a popular text book on radiolarians presented by De Wever et al. (2001). The zonal scheme has been widely utilized for the dating of marine sequences accumulated in the Panthalassa and Tethys (e.g., Matsuoka et al., 1998, 2002; Zamoras and Matsuoka, 2001, 2004; Li et al., 2016, 2018; Ito and Matsuoka, 2018).

Radiolarian zonation

As stated in "Introduction", much effort has been made on taxonomic and biostratigraphic research for Jurassic radiolarians. It is necessary to update zonal names and definition in accordance with the progress in radiolarian research for the last two decades. The updated

zonal scheme for the Jurassic in Japan and the Pacific is summarized in Fig. 1. Vertical distributions of selected radiolarian species, especially zone-diagnostic taxa, are also illustrated in Fig. 1.

To define zones, radiolarian bio-events such as the evolutionary first appearance biohorizon (EFAB), first occurrence biohorizon (FOB), and last occurrence biohorizon (LOB) were used. The EFAB, which is the most synchronously reliable among various kinds of biohorizons, is selected as often as possible.

Bipedis horiae Zone

Code. JR0

Base. LOB of *Haeckelicyrtium breviora* Sugiyama.

Top. FOB of *Parahsuum simplum* Yao.

Remarks. Sugiyama (1997) established the *Haeckelicyrtium breviora*–*Bipedis horiae* Partial-range Zone (JR0A) and the *Bipedis horiae* Lowest occurrence Zone (JR0B) in the Sakahogi section of the Mino Terrane, central Japan. The *Bipedis horiae* Zone (JR0) of the present study is a zonal unit for the lowest portion of the Jurassic, combining JR0A and JR0B proposed by Sugiyama (1997). The LOB of *H. breviora* Sugiyama corresponds to the Triassic–Jurassic boundary (TJB) in the Sakahogi section (Sugiyama, 1997). Since species belonging to the genus *Bipedes* are common in marine sequences around the TJB, the phyletic analysis of *Bipedis* species can contribute to further refinement of this zone. The first occurrence of *B. horiae* Sugiyama has been reported only in the Sakahogi section so far, which is located just above the TJB.

Age. Hettangian to early Sinemurian.

Parahsuum simplum Zone

Code. JR1

Base. FOB of *Parahsuum simplum* Yao.

Top. FOB of *Trillus elkhornensis* Pessagno and Blome.

Remarks. Sugiyama (1997) showed the first occurrence of *Parahsuum simplum* Yao in the Sakahogi section, being higher than the FOB of *B. horiae*. The first occurrence biohorizon of *P. simplum* is located within the Sinemurian in Haida Gwaii, Canada (Carter et al., 1998).

Age. Late Sinemurian to middle Early Pliensbachian.

Trillus elkhornensis Zone

Code. JR2

Base. FOB of *Trillus elkhornensis* Pessagno and Blome.

Top. FOB of “*Laxtorum*” *jurassicum* Isozaki and Matsuda.

Remarks. Carter et al. (2010) reported that *Trillus elkhornensis* Pessagno and Blome first




Fig. 1. Updated radiolarian zonation for the Jurassic in Japan and the western Pacific. Chronostratigraphy is based on Ogg et al. (2016). Images of index radiolarians are reprinted from authors' previous papers (Matsuoka, 1983, 1998, 2004) and unpublished data. Arrows in the range chart indicate that the vertical ranges extend beyond the Jurassic-Cretaceous boundary.

occurs in the *Hsuum mulleri*-*Trillus elkhornensis* Zone of middle Lower Pliensbachian.
Age. Middle Early Pliensbachian to Toarcian.

"Laxtorum" jurassicum Zone

Code. JR3

Base. FOB of *"Laxtorum" jurassicum* Isozaki and Matsuda.

Top. EFAB of *Striatojaponocapsa plicarum* (Yao).

Age. Aalenian.

Striatojaponocapsa plicarum Zone

Code. JR4

Base. EFAB of *Striatojaponocapsa plicarum* (Yao).

Top. EFAB of *Striatojaponocapsa conexa* (Matsuoka).

Remarks. *Striatojaponocapsa* was established by Kozur (1984), taking *Tricolocapsa plicarum* Yao as the type species. The evolutionary lineage from the ancestor *S. plicarum* to the descendant *S. conexa* was first demonstrated by Matsuoka (1983). O'Dogherty et al. (2006) described two new species of *Striatojaponocapsa* in the Austrian Alps; *S. synconexa* O'Dogherty, Goričan, and Dumitrica and *S. riri* O'Dogherty, Goričan, and Dumitrica. Hatakeda et al. (2007) examined in detail the evolutionary history of the *Striatojaponocapsa* lineage in continuous sections in Japan. They concluded that *S. plicarum* continuously and gradually changed into *S. synconexa*, then *S. synconexa* was rapidly replaced by *S. conexa* in a phyletic evolutionary process. In accordance with the change of generic assignment of the nominal species, the zonal name is changed to the *Striatojaponocapsa plicarum* Zone.

Age. Bajocian to early Bathonian.

Striatojaponocapsa conexa Zone

Code. JR5

Base. EFAB of *Striatojaponocapsa conexa* (Matsuoka).

Top. EFAB of *Kilinora spiralis* (Matsuoka).

Remarks. In relation to the change of generic assignment of the nominal species (Hull, 1997), the zonal name is changed to the *Striatojaponocapsa conexa* Zone.

Age. Late Bathonian to middle Callovian.

Kilinora spiralis Zone

Code. JR6

Base. EFAB of *Kilinora spiralis* (Matsuoka).

Top. LOB of *Striatojaponocapsa conexa* (Matsuoka).

Remarks. The genus *Kilinora* was established by Hull (1997) who took *Stylocapsa(?) spiralis*

Matsuoka as the type species. In accordance with the change of generic assignment of the nominal species, the zonal name is changed to the *Kilinora spiralis* Zone. The evolutionary lineage form the ancestor *Kilinora hemicostata* (Matsuoka) to the descendant *K. Spiralis* was first reported at several land sections in Japan (Matsuoka, 1983, 1995a). The same lineage was recognized at DSDP Site 534 in the north-western Central Atlantic as well (Matsuoka and Baumgartner, 1997).

Age. Late Callovian to Oxfordian.

Hsuum maxwelli Zone

Code. JR7

Base. LOB of *Striatojaponocapsa conexa* (Matsuoka).

Top. LOB of *Hsuum maxwelli* Pessagno.

Age. Kimmeridgian.

Loopus primitivus Zone

Code. JR8

Base. LOB of *Hsuum maxwelli* Pessagno.

Top. EFAB of *Pseudodictyomitra carpatica* Lozyniak.

Remarks. Yang (1993) established the genus *Loopus*, taking *Pseudodictyomitra primitiva* Matsuoka and Yao as the type species. In relation to the change of generic assignment of the nominal species, the zonal name is changed to the *Loopus primitivus* Zone. The evolutionary first appearance of *Pseudodictyomitra carpatica* was reported at Site 801 of ODP Leg 129 (Yang and Matsuoka, 1997; Matsuoka and Yang, 2000). Zügel (1997) showed the occurrence of *P. carpatica* in the lower Tithonian of the Solnhofen area, southern Germany.

Age. Early Tithonian.

Pseudodictyomitra carpatica Zone

Code. KR1

Base. EFAB of *Pseudodictyomitra carpatica* (Lozyniak).

Top. EFAB of *Cecrops septemporatus* (Parona).

Remarks. The Jurassic–Cretaceous boundary is located within this zone.

Age. Late Tithonian to early Valanginian.

Concluding remarks

The current status and future directions in Mesozoic radiolarian biochronological research are recently documented by Goričan et al. (2018). They stress the importance of

detailed documentation of evolutionary first and last appearance in different radiolarian phylogenetic lineages. In the last two decades, some new taxa have been proposed and their biostratigraphic significance has been discussed. On the basis of the progress, this article updated Jurassic radiolarian zonation.

Generic assignment of many species, even zone-diagnostic species (i.e., "*Laxtorum jurassicum*"), is still debatable. Furthermore, phylogenetic relationships among Early and Late Jurassic radiolarians are not entirely clear compared to those of Middle Jurassic radiolarians. We expect this article to act as a stepping stone to further progress.

Acknowledgements

We thank Peter O. Baumgartner, Špela Gorican, Luis O'Dogherty and Elizabeth S. Carter for long lasting discussion on Jurassic radiolarian biostratigraphy. Toshiyuki Kurihara reviewed the manuscript and gave us constructive comments. Theodore Brown read the manuscript and corrected the English. This work was financially supported by JSPS KAKENHI Grant Numbers 15K05329 and 15H02142 to A. Matsuoka.

References

- Carter, E. S., Whalen, P. A. and Guex, J., 1998, Biochronology and paleontology of Lower Jurassic (Hettangian and Sinemurian) radiolarians, Queen Charlotte Islands, British Columbia. *Geol. Surv. Canada, Bull.*, **496**, 1–162.
- Carter, E. S., Gorican, Š., Guex, J., O'Dogherty, L., De Wever, P., Dumitrica, P., Hori, R. S., Matsuoka, A. and Whalen, P. A., 2010, Global radiolarian zonation for the Pliensbachian, Toarcian and Aalenian. *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, **297**, 401–419.
- De Wever, P., Dumitrica, P., Caulet, J. P. and Caridroit, M., 2001, Radiolarians in the sedimentary record. G.B.S. Publishers, Amsterdam, 533 p.
- Gorican, Š., Carter, E. S., Dumitrica, P., Whalen, P. A., Hori, R. S., De Wever, P., O'Dogherty, L., Matsuoka, A. and Guex, J., 2006, *Catalogue and systematics of Pliensbachian, Toarcian and Aalenian radiolarian genera and species*. Založba ZRC/ZRC Publishing, ZRC SAZU, Ljubljana, 446 p.
- Gorican, Š., Carter, E. S., Guex, J., O'Dogherty, L., De Wever, P., Dumitrica, P., Hori, R. S., Matsuoka, A. and Whalen, P. A., 2013, Evolutionary patterns and palaeobiogeography of Pliensbachian and Toarcian (Early Jurassic) Radiolaria. *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, **386**, 620–636.
- Gorican, Š., O'Dogherty, L., Baumgartner, P. O., Carter, E. S., Matsuoka, A., 2018, Mesozoic radiolarian biochronology – current status and future directions. *Rev. Micropaléont.*, **61**, 165–189.
- Hatakeda, K., Suzuki, N. and Matsuoka, A., 2007, Quantitative morphological analyses of evolutionary history of the Middle Jurassic polycystine radiolarian genus *Striatojaponocapsa* Kozur. *Marine Micropaleont.*, **63**, 39–56.
- Hori, N., 1999, Latest Jurassic radiolarians from the northeastern part of the Torinoko Block, Yamizo Mountains, central Japan. *Sci. Rep. Inst. Geosc., Univ. Tsukuba, Sec. B (Geol. Sci.)*, no. 20, 47–114.
- Hull, D. M., 1997, Upper Jurassic Tethyan and southern Boreal radiolarians from western North America. *Micropaleontology*, **43**, 1–202.
- Ito, T. and Matsuoka, A., 2018, Lithology and radiolarian age of the Ryokami-yama Chert Formation in eastern Mt. Ryokami: Possible décollement zone in Permian pelagic sequence in mid-Mesozoic accretionary

- complexes of Southwest Japan. *Isl. Arc*, **27**, e12273.
- Kiessling, W., 1999, Late Jurassic radiolarians from the Antarctic Peninsula. *Micropaleontology*, **45**, 1–96.
- Kozur, H., 1984, New radiolarian taxa from the Triassic and Jurassic. *Geologisch Paläontologische Mitteilungen, Innsbruck*, **13**, 49–88.
- Li, X., Matsuoka, A., Li, Y. L., Wei, Y. S. and Wang, C. S., 2016, Middle Jurassic to Early Cretaceous radiolarian assemblages from the chert blocks in the mélange along the Yarlung–Tsangpo Suture Zone, near Zhongba County, Southern Tibet. *News Osaka Micropaleontol. (NOM), Spec. Vol.*, no. 16, 79–94.
- Li, X., Matsuoka, A., Li, Y. L., Wei, Y. S. and Wang, C. S., 2018, Radiolarian-based study on the fabric and the formation process of the Early Cretaceous mélange near Zhongba, Yarlung–Tsangpo Suture Zone, southern Tibet. *Isl. Arc*, **27**, e12282.
- Matsuoka, A., 1983, Middle and Late Jurassic radiolarian biostratigraphy in the Sakawa and adjacent areas, Shikoku, Southwest Japan. *J. Geosci. Osaka City Univ.*, **26**, 1–48.
- Matsuoka, A., 1992, Jurassic and Early Cretaceous radiolaria from the western Pacific: Leg 129 of the Ocean Drilling Program. In Larson R., Lancelot Y., et al, eds., *Proc. Ocean Drill. Program, Sci. Results*, **129**, 203–220.
- Matsuoka, A., 1995a, Jurassic and Lower Cretaceous radiolarian zonation in Japan and in the western Pacific. *Isl. Arc*, **4**, 140–153.
- Matsuoka, A., 1995b, Middle Jurassic to Early Cretaceous radiolarian occurrences in Japan and the western Pacific (ODP Sites 800-801). In Baumgartner, P. O., O'Dogherty, L., Goričan, Š., Urquhart, E., Pillecuit, A. and De Wever, P., eds., Middle Jurassic to Lower Cretaceous Radiolaria of Tethys: Occurrences, Systematics, Biochronology. *Mém. Géol. (Lausanne)*, no. 23, 937–966.
- Matsuoka, A., 1998, Faunal composition of earliest Cretaceous (Berriasian) radiolaria from the Mariana Trench in the western Pacific. *News Osaka Micropaleontol. (NOM), Spec. Vol.*, no. 11, 165–187.
- Matsuoka, A., 2004, Toarcian (Early Jurassic) radiolarian fauna from the Nanjo Massif in the Mino terrane, central Japan. *News Osaka Micropaleontol. (NOM), Spec. Vol.*, no. 13, 69–87.
- Matsuoka, A., 2006, Revised radiolarian zonation for the entire Jurassic in the western Pacific and Japan. *Volumina Jurassica*, **4**, 183–184.
- Matsuoka, A. and Baumgartner, P. O., 1997, Middle Jurassic radiolarians from the basal sediments at DSDP Site 534, Blake Bahama Basin, Northern Atlantic. *News Osaka Micropaleontol. (NOM), Spec. Vol.*, no. 10, 183–191.
- Matsuoka, A., Yamakita, S., Sakakibara, M. and Hisada, K., 1998, Unit division for the Chichibu Composite Belt from a view point of accretionary tectonics and geology of western Shikoku, Japan. *J. Geol. Soc. Japan*, **104**, 634–653 (in Japanese with English abstract).
- Matsuoka, A. and Yang, Q., 2000, A direct correlation between North American and Japan-Pacific radiolarian zonal schemes for the Upper Jurassic. *GeoResearch Forum*, **6**, 119–128.
- Matsuoka, A. and Yao, A., 1986, A newly proposed radiolarian zonation for the Jurassic of Japan. *Marine Micropaleont.*, **11**, 91–106.
- Matsuoka, A., Yang, Q., Kobayashi, K., Takei, M., Nagahashi, T., Zeng, Q. G. and Wang, Y. J., 2002, Jurassic–Cretaceous radiolarian biostratigraphy and sedimentary environments of the Cenozoic Tethys: records from the Xialu Chert in the Yarlung–Zangbo Suture Zone, southern Tibet. *J. Asian Earth Sci.*, **20**, 277–287.
- O'Dogherty, L., Bill, M., Goričan, Š., Dumitrica, P. and Masson, H., 2006, Bathonian radiolarians from an ophiolitic melange of the Alpine Tethys (Gets Nappe, Swiss-French Alps). *Micropaleontology*, **51**, 425–485.
- O'Dogherty, L., Carter, E. S., Dumitrica, P., Goričan, Š., De Wever, P., Bandini, A. N., Baumgartner, P. O. and Matsuoka, A., 2009, Catalogue of Mesozoic radiolarian genera. Part 2: Jurassic–Cretaceous. *Geodiversitas*, **31**, 271–356.
- O'Dogherty, L., Goričan, Š. and Gawlick, H. J., 2017, Middle and Late Jurassic radiolarians from the Neotethys suture in the Eastern Alps. *J. Paleont.*, **91**, 25–72.
- Ogg, J. G., Ogg, G. M. and Gradstein, F. M., 2016, *A Concise Geologic Time Scale 2016*. Elsevier, Amsterdam, 234 p.
- Pessagno, E. A. Jr., Cantú-Chapa, A., Mattinson, J. M., Meng, X. and Karimnia, S. M., 2009, The Jurassic–Cretaceous boundary: new data from North America and the Caribbean. *Stratigraphy*, **6**, 185–262.
- Steiger, T., 1992, Systematik, stratigraphie und Palökologie der Radiolarien des Oberjura-Unterkreiden-

- Grenzbereiches im Osterhorn-Tirolikum (Nördliche Kalkalpen, Salzburg und Bayern. *Zitteliana*, **19**, 3–188 (in German with English abstract).
- Sugiyama, K., 1997, Triassic and Lower Jurassic radiolarian biostratigraphy in the siliceous claystone and bedded chert units of the southeastern Mino Terrane, Central Japan. *Bull. Mizunami Fossil Mus.*, no. 24, 79–193.
- Yang, Q., 1993, Taxonomic studies of Upper Jurassic (Tithonian) radiolaria from the Taman Formation, east-central Mexico. *Palaeoworld*, **3**, 1–164.
- Yang, Q. and Matsuoka, A., 1997, A comparative study on Upper Jurassic radiolarian biostratigraphy of the Taman Formation, east-central Mexico and the ODP Site 801B Section, west Pacific. *Sci. Rep. Niigata Univ. (Geol.)*, no. 12, 29–49.
- Yeh, K.-Y., 2009, A middle Jurassic radiolarian fauna from South Fork Member of Snowshoe Formation, east-central Oregon. *Coll. Res., Nati. Mus. Natu. Sci., Taiwan*, **22**, 15–125.
- Zamoras, L. R. and Matsuoka, A., 2001, Malampaya Sound Group: a Jurassic-Early Cretaceous accretionary complex in Busuanga Island, North Palawan Block (Philippines). *J. Geol. Soc. Japan*, **107**, 316–336.
- Zamoras, L. R. and Matsuoka, A., 2004, Accretion and postaccretion tectonics of the Calamian Islands, North Palawan block, Philippines. *Isl. Arc*, **13**, 506–519.
- Zügel, P., 1997, Discovery of a radiolarian fauna from the Tithonian of the Solnhofen area (Southern Franconian Alb, southern Germany). *Palaeontologische Zeitschrift*, **71**, 197–209.