

Carboniferous brachiopod *Latiproductus edelburgensis* (Phillips, 1836) from Akiyoshi and Omi, Japan

Yousuke IBARAKI*, Yasuhiro OTA** and Masayuki FUJIKAWA***

Abstract

A productid brachiopod species, *Latiproductus edelburgensis* (Phillips), is described from the upper Viséan to lower Serpukhovian of the Akiyoshi Limestone, Akiyoshi, southwest Japan and the Omi Limestone, Omi, central Japan. The occurrence of *L. edelburgensis* indicates a late Viséan to early Serpukhovian age for the *Mediocris mediocris* Zone of the Akiyoshi Limestone and the *Eostaffella-Millerella* Zone of the Omi Limestone. The stratigraphic and geographic distributions of *L. edelburgensis* are restricted to the Lower Carboniferous (upper Viséan to lower Serpukhovian) of Europe, central and eastern Asia, northern Africa and western Panthalassa, and completely absent in North and South America and Australia.

Key words: Akiyoshi, Brachiopoda, *Latiproductus edelburgensis*, lower Carboniferous, Omi.

Introduction

Gigantoproductoids are large-sized brachiopods known as a leading fossil of Early Carboniferous (late Viséan–early Serpukhovian) from Europe, Algeria, Russia, China and North America. In Japan, several gigantoproductoid species were previously described from the Omi Limestone in Omi, Itoigawa City, central Japan (Hayasaka, 1924), the Akiyoshi Limestone in Akiyoshi Plateau, southwest Japan (Yanagida, 1979), the Koyama Limestone in Oga, Takahashi City, southwest Japan (Ibaraki et al., 2014), the Ichinotani Formation in the Fukuji area, Hida Mountains (Tazawa and Kato, 1986; Ibaraki et al., 2009), the Tsuchikurazawa

* Fossa Magna Museum, Itoigawa 941-0056, Japan

** Kitakyushu Museum of Natural History and Human History, Kitakyushu 805-0071, Japan

*** Akiyoshidai Science Museum, Mine 754-0511, Japan

Corresponding author: Yousuke Ibaraki,

yousuke.ibaraki@city.itoigawa.lg.jp

(Manuscript received 1 March, 2021; accepted 20 March, 2021)




Fig. 1. Map showing the fossil localities, Loc. 1 (Okubo), Loc. 2 (Nishiyama Quarry), A: Akiyoshi, B: Omi, using the topographical map of “Akiyoshidaihokubu”, “Akiyoshidai” and “Itoigawa” scale 1:25,000 published by the Geographical Survey Institute of Japan.

Limestone in Kotaki, Itoigawa City, central Japan (Tazawa, 2004; Ibaraki et al., 2008, 2010) and the Onimaru Formation in the Nagaiwa-Onimaru area and the Hikoroichi Formation in the Hikoroichi area, southern Kitakami Mountains, northeast Japan (Tazawa and Miyake, 2002; Tazawa and Ibaraki, 2009; Tazawa, 2018; Tazawa and Taira, 2020).

In this paper, we examine four specimens of gigantoproductoids from the *Mediocris mediocris* Zone of the Akiyoshi Limestone in Shishidedai and Okubo, Akiyoshi Plateau, southwest Japan and the *Eostaffella-Millerella* Zone of the Omi Limestone in Nishiyama Quarry, Omi, Itoigawa City, central Japan (Fig. 1).

Material

The Akiyoshi specimens were collected by late Professor Juichi Yanagida of Kyushu University. Most of the Omi specimens were collected by Emeritus Professor Jun-ichi Tazawa of Niigata University. Based on the present systematic descriptions, all of the specimens are referred to *Latiproductus edelburgensis* (Phillips, 1836), originally described from the Lower Carboniferous of Bolland, Lancashire, England, the UK.

The specimens described herein are registered and housed in the Fossa Magna Museum



Fig. 2. Geographical distribution of *Latiproductus edelburgensis* (Phillips) in the late Viséan (adapted Scotese, 2004). 1: Algeria, 2: Lancashire, 3: Yorkshire, 4: Pennine Mountains, 5: Silesia, 6: Holy Cross Mountains, 7: Moscow Basin, 8: northern Timan, 9: Pechora, 10: central Ural, 11: central Kazakhstan, 12: northern Kirgizia, 13: Tien-Shan, 14: Xizang, 15: Qinghai, 16: Jilin, 17: Sichuan, 18: Jiangxi, 19: Hubei, 20: Hunan, 21: Guizhou, 22: Akiyoshi, 23: Omi, 24: Oga, 25: Nagaiwa-Onimaru.

(specimen numbers prefixed with FMM) and the Kitakyushu Museum of Natural History and Human History (specimen numbers prefixed with KMNHIVP).

Stratigraphic and geographic distributions of *Latiproductus edelburgensis*

Latiproductus edelburgensis has been described from the upper Viséan–Serpukhovian of Algeria (1; Legrand-Brain, 1973, 1980; Pareyn, 1961). Lancashire, England (2; Phillips, 1836), Yorkshire, England (3; Davidson, 1958–1963; Prentice, 1956), Pennine Mountains, England (4; Pattison, 1981), Silesia, Germany (5; Paeckelmann, 1931), Holy Cross Mountains, Poland (6; Zakowa, 1986), Moscow Basin, western Russia (7; Sarytcheva and Sokolskaya, 1952), northern Timan, western Russia (8; Aisenberg and Poletaev, 1970), Pechora, western Russia (9; Kalashnikov, 1974), central Urals, western Russia (10; Einor, 1957), central Kazakhstan (11; Litovinovich, Aksenova and Razina, 1969), northern Kirgizia (12; Gladchenko, 1955; Galitzkaja, 1977), Tien-Shan, Northwest China (13; Gröber, P., 1909; Krenkel, 1913), Xizang, Southwest China (14; Jin et al., 1985), Qinghai, Northwest China (15; Yang et al., 1962), Jilin, Northeast China (16; Lee et al., 1980), Sichuan, Southwest China (17; Yang and Jiang, 1987), Jiangxi, South China, (18; Wang et al., 1982), Hubei, South China (19; Wang, 1984), Hunan, South China (20; Tan, 1987; Liu et al., 1982), Guizhou, South China (21; Chao, 1927), Akiyoshi, southwest Japan (22; Yanagida, 1989), Omi, central Japan (23; This paper), Oga, southwest Japan (24; Ibaraki et al., 2014), Nagaiwa-Onimaru, South Kitakami Mountains, northeast Japan (25; Tazawa and




Fig. 3. A₁ – A₄: ventral, posterior, lateral views and enlarged radial costae of conjoined valve, KMNHiv710001. B₁ – B₄: ventral, posterior, lateral views and enlarged radial costae of dorsal valve, KMNHiv710002. Scale bars are 1 cm, except of A₄ and B₄.

Taira, 2020) (Fig. 2).

From the above data, the range of *Latiproductus edelburgensis* is assigned to late Viséan–Serpukhovian, which is the same as the summary by Brunton et al. (2000). Qiao and Shen (2015) also concluded as above. Palaeobiogeographically, it is noteworthy that *L. edelburgensis* has been mostly found from equatorial to mid latitude areas of Asia and Europe, but completely absent in North America, South America and Australia. This result is consistent with one of the conclusions in Qiao and Shen (2014). The lower Carboniferous

limestones of Omi and Akiyoshi were probably reef-seamount of the Panthalassa in the mid-latitude area of the Northern Hemisphere during late Viséan–Serpukhovian. This conclusion is consistent with that of Tazawa et al. (2005), in which they noted that the Akiyoshi-Omi reef-seamounts were probably located at the lower to middle northern palaeolatitude on the Panthalassa during the early Carboniferous (late Viséan).

Systematic descriptions

(by Y. Ibaraki)

Order Procutida Sarycheva and Sokolskaya, 1959
 Suborder Productidina Waagen, 1883
 Superfamily Linoproductoidea Stehli, 1954
 Family Linoproductidae Stehli, 1954
 Subfamily Gigantoproductinae Muir-Wood and Cooper, 1960
 Tribe Semiplanini Sarycheva, 1960
 Genus *Latiproductus* Sarycheva and Legrand-Blain, 1977

Type species.—*Productus latissimus* Sowerby, 1822.

Latiproductus edelburgensis (Phillips, 1836)

Figs. 3–4

Producta edelburgensis Phillips, 1836, p. 214, pl. 7, fig. 5.

Productus giganteus mut. *edelburgensis* (Phillips). Glöber, 1909, p. 372, pl. 1, fig. 11; pl. 2, figs. 3–4.

Productus giganteus var. *edelburgensis* (Phillips). Hayasaka, 1924, p. 143, pl. 54, figs. 1, 2.

Striatifera edelburgensis (Phillips). Chao, 1927, p. 107, pl. 10, figs. 4, 5; pl. 12, fig. 6.

Gigantella edelburgensis var. *glöbereri* Sarycheva, 1928, p. 51, pl. 4, fig. 5.

Productus (*Gigantella*)? *edelburgensis* (Phillips). Paeckelmann, 1931, p. 260, pl. 29, figs. 1a–1c.

Productus (*Gigantella*) *edelburgensis*? Ozaki, 1939, p. 244, pl. 38, fig. 2.

Productus (*Gigantella*) *edelburgensis*? var. *shajenwaensis* Ozaki, 1939, p. 242, pl. 37, fig. 4; pl. 38, fig. 1; pl. 39, fig. 2.

Gigantoproductus edelburgensis (Phillips). Sarycheva in Sarycheva and Sokolskaya, 1952, p. 131, pl. 35, fig. 180; Prentice, 1956, p. 234, pl. 20, figs. 1a–c, 2; Galitskaya, 1977, p. 147, pl. 61, figs. 2a–2b; pl. 62, fig. 1; pl. 63, figs. 1–2.

Productus (*Gigantoproductus*) *edelburgensis* (Phillips). Gladchenko, 1955, p. 19, pl. 10, figs. 1a–1c.

Productus (*Gigantoproductus*) *edelburgensis* var. *schaitankaensis* Einor, 1957, p. 151, pl. 3, fig. 3.

Productus-Gigantoproductus-edelburgensis (Phillips). Pareyn, 1961, p. 202, pl. 24, figs. 4, 5.




Fig. 4. A1 – A4: ventral, posterior, lateral views and enlarged radial costae of ventral valve, FMM2023. B1 – B4: ventral, posterior, lateral views and enlarged radial costae of ventral valve, FMM2024. Scale bars are 1 cm, except of A4 and B4.

Productus (Gigantoproductus) edelburgensis var. *beleutensis* Litvinovich, 1962, p. 207, pl. 7, fig. 2.

Gigantoproductus cf. *edelburgensis* (Phillips). Legrand-Blain, 1973, p. 91, pl. 2, figs. 1a–1b.

Gigantoproductus edelburgensis edelburgensis (Phillips). Semichatova, 1975, p. 173, pl. 77, fig. 1.

Gigantoproductus edelburgensis schaitakaensis (Einor). Semichatova, 1975, p. 173, pl. 77, fig. 2.

Gigantoproductus edelburgensis? var. *shajenwaensis* (Ozaki). Yang et al., 1977, p. 368, pl. 146, fig. 4.

Gigantoproductus aff. *edelburgensis* (Phillips). Yanagida, 1979, p. 112, fig. 3.

Gigantoproductus aff. *submaximus* (Bolikhovitinova). Yanagida, 1979, p. 111, fig. 2.

Gigantoproductus edelburgensis var. *shajenwaensis* (Ozaki). Liu et al., 1982, p. 189; pl. 135, fig. 5.

Latiproductus (?) *edelburgensis sahariensis* forme a Legrand-Blain, 1980, p. 42, pl. 3, figs. 3; text-figs. 13, 15.

Latiproductus (?) *edelburgensis sahariensis* forme c Legrand-Blain, 1980, p. 46, pl. 5, figs. 1, 2; text-figs. 13, 16.

Latiproductus (?) *edelburgensis subsahariensis*. Legrand-Blain, 1980, p. 47, pl. 1, fig. 6; pl. 3, fig. 4; text-figs. 14, 17.

Latiproductus edelburgensis (Phillips). Zakowa, 1986, p. 65, pl. 6, fig. 1; Ibaraki et al., 2014, p. 17, fig. 3; Tazawa and Taira, 2020, p. 21, fig. 7.

Material.—(1) An imperfect conjoined valve, KMNHIv710001 from Loc. 1; (2) two ventral valves, FMM2023, 2024 from Loc. 2; (3) a dorsal valve, KMNHIv710002 from Loc. 1.

Description.—Shell medium size for genus, transversely semicircular in outline, with greatest width at hinge; length 60 mm, width 110 mm in the largest specimen (FMM2023); length 50 mm, width about 90 mm in the smallest specimen (FMM2024). Ventral valve moderately convex in lateral profile, strongly geniculated at 40–50 mm from umbo; flanks gently inclined; umbo small, rounded and inflated; ears large, triangular and moderately demarcated from flanks; sulcus absent. External surface of ventral valve ornamented with numerous costae but no rugae; costae regular in anterior part, but irregular in trail, numbering 8–10 per 10 mm at about midvalve, and costae sometimes bifurcated or inserted in anterior regions; intercostal sulci as wide as costae; several radial fluting crossing costae on trail; numerous fine growth lines on valve; spines or spine bases not preserved on the surface of valves. Dorsal valve slightly concave, geniculated at 40–50 mm from umbo; incurve of midvalve weaker than that of ventral valve; external surface of dorsal disc ornamented with numerous costae; costae regular, numbering 8–9 per 10 mm at about midvalve; costae sometimes bifurcated or inserted in anterior regions; no rugae; numerous fine growth lines over the inner surface; interior of both valves not observed.

Remarks.—The specimens are assigned to *Latiproductus edelburgensis* (Phillips, 1836), originally described from the Lower Carboniferous of Bolland, Lancashire, England, from account of size, shape, and external ornament of ventral valve, particularly in its size, shape of umbo. *Latiproductus latissimus* (Sowerby, 1822), described and figured by Pattison (1981) differs from *L. edelburgensis* in its smaller size and flattened ears, existence of a sulcus, finer costae, absence of intercostal sulci on ventral valve and thicker shell. *Gigantoproductus submaximus* (Bolikhovitinova, 1932) resembles *L. edelburgensis* in shape of ventral valve and

the presence of several rugae on trail, but differs in its larger size, finer costae on ventral valve. A specimen described as *Gigantoproductus* aff. *submaximus* (Bolkhovitnova) by Yanagida (1979) is reassigned to *Latiproductus edelburgensis*, owing to the following reasons: 1) The specimen of Yanagida (1979) lacks a long trail with fluting flexuously costae in the ventral valve, which are characteristics of *L. edelburgensis* as shown in well-preserved specimen (Galitskaya, 1977: fig. 1, pl. 63) and 2) the ventral valve of *G. submaximus* is much larger and having finer costae than those of *L. edelburgensis*.

Acknowledgements

The first author extends cordially thank Jun-ichi Tazawa, Professor Emeritus Niigata University, for his earnest guidance and encouragement throughout the present study. We thank anonymous reviewers for critically reading the manuscript, by which this paper was greatly improved.

References

- Aisenberg, D. E. and Poletaev, V. I., 1970, Brachiopoda. In Barkhatova, V. P., ed., *Carboniferous and Lower Permian biostratigraphy of northern Timan*. Nedra, Leningrad, 3–228 (in Russian).
- Bolkhovitnova, M. A., 1932, New information of stratigraphy of Eugo-east part, general geological map, 58th sheet, Nauchotek (ed.), Moskva, 49 p. (in Russian).
- Brunton, C. H. C., Lazarev, S. S., Grant, R. E. and Jin, Y. G., 2000., Productidina. In Kaesler, R. L., ed., *Treatise on Invertebrate Paleontology, Part H Brachiopoda Revised, Volume 3: Linguliformea, Craniiformea and Rhynchonelliformea (Part)*, Geol. Soc. Amer., Boulder and Univ. Kansas, Lawrence, 424–609.
- Chao, Y. T., 1927, Productidae of China. Part 1. Producti, *Palaeont. Sinica, Ser. B*, Fasc. 2, 83–120.
- Davidson, T., 1858–1863, *British fossil Brachiopoda, Vol. 2. Permian and Carboniferous species*. Palaeontograph. Soc., London, 280 p.
- Einor, O. L., 1957, Carboniferous brachiopod fauna from the western slope in Central Ural. *Ezh. Vsesoy. Paleont. Obsch.*, **16**, 142–161 (in Russian).
- Galitskaya, A. Ya, 1977, *Early and Middle Carboniferous Productids from Northern Kyrgyzstan*. Ilim, Frunze, 297 p. (in Russian).
- Gladchenko, A. Ya, 1955, *Field atlas of index fossils of Brachiopods from the lower Carboniferous of northern Kirgizia*. Akad. Nauk Kirgiz., SSR, Frunze, 30 p. (in Russian).
- Glöber, P., 1909, Carbon und Carbonfossilien des nördlichen und zentralen Tian-Schan. *Abh. König. Akad. Wissenschaft. 2. Klas.*, **24**, 341–384.
- Hayasaka, I., 1924, On the fauna of anthracolithic limestone of Omi-mura in the western part of Echigo. *Sci. Rep. Tohoku Imp. Univ., 2nd Ser.*, **8**, 1–83.
- Ibaraki, Y., Tazawa, J., Sato, K. and Nakamura, Y., 2008, *Gigantoproductus* (Carboniferous Brachiopoda) from Kotaki, Itoigawa City, Niigata Prefecture, central Japan. *Sci. Rep. Niigata Univ., (Geol.)*, no. 23, 55–64.
- Ibaraki, Y., Tazawa, J. and Nakamura, Y., 2010, Additional *Gigantoproductus* species from the upper Viséan–Namurian limestone of Kotaki, central Japan. *Sci. Rep. Niigata Univ., (Geol.)*, no. 25, 63–68.
- Ibaraki, Y., Miyake, Y. and Tazawa, J., 2014, Early Carboniferous (late Viséan) brachiopods from the Koyama Limestone of Kamiotake in the Oga area, Okayama Prefecture, Southwest Japan. *Earth Sci. (Chikyū Kagaku)*, **68**, 69–79.
- Ibaraki, Y., Tazawa, J. and Miyake, Y., 2009, *Gigantoproductus* (Carboniferous Brachiopoda) from the lowest part of the Ichinotani Formation, Fukui, Hida Gaien Belt, central Japan. *Sci. Rep. Niigata Univ., (Geol.)*, no. 24, 1–5.

- Jin, Y., Wang, Y., Sun, D. and Shi, Q., 1985, Late Paleozoic and Triassic brachiopods from the East of the Quighan-Xizang Plateau. In Regional Geological Surveying Team of Sichuan Province and Nanjing Institute of Geology and Palaeontology, eds., *Stratigraphy and Palaeontology in W. Sichuan and E. Xizang, China (Part 3)*. Sichuan Sci. Tec. Press, Chengdu, 182-249 (in Chinese).
- Kalashnikov, N. V., 1974, *Early Carboniferous brachiopod from Pechora Urals*. Nauka, Leningrad, 220 p. (in Russian).
- Krenkel, E., 1913, Wissenschaftliche ergebnisse der Reise von Prof. Dr. G. Merzbacher im zentralen und östlichen Tian-Schan 1907/8. *Abh. Königlich Bayerischen Akad. Wissen. Math. -phys.*, klasse 3, band. 8, 1-44.
- Lee, L., Gu, F. and Su, Y., 1980, Carboniferous and Permian Brachiopoda. In Shenyang Institute of Geology and Mineral Resources eds., *Paleontological Atlas of Northeast China, Paleozoic Volume*, Geol. Pub. House, Beijing, 327-428 (in Chinese).
- Legrand-Blain, M., 1973, Les gigantoproductides (brachiopodes) du Sahara Algerien. *Bull. Soc. Hist. Natur. de l'Afrique du Nord*, **64**, 79-157.
- Legrand-Blain, M., 1980, Les gigantoproductides (brachiopodes) du Sahara Algerien 3-Semiplanidae Viseens et Namuriens. *Bull. Soc. Hist. Natur. de l'Afrique du Nord*, **69**, 3-85.
- Litvinovich, N. V., 1962, Carboniferous and Permian Deposits of the Western Part of Central Kazakhstan. *Izd. Moskov. Univ., Moskva*, 389 p. (in Russian).
- Litvinovich, N. V., Aksenova, G. G. and Razina, T. P., 1969, *Stratigraphy and lithology of the Lower to Upper Carboniferous in central western part of Kazakhstan*. Nedra, Moskva, 447 p. (in Russian).
- Liu, Z., Tan, Z. and Ding, Y., 1982, Brachiopoda. In Geological Bureau of Hunan eds., *Palaeontological Atlas of Hunan Province*. Geol. Pub. House, Beijing, 172-216 (in Chinese).
- Muir-Wood, H. M. and Cooper, G. A., 1960, Morphology, classification and life habits of the Productoidea (Brachiopoda). *Geol. Soc. America Mem.*, **81**, 447 p.
- Ozaki, K., 1939, On some Lower Carboniferous brachiopods from central Hunan, China. *Shanghai Sci. Inst. Jour. Section 2*, **2**, 225-282.
- Paeckelmann, W., 1931, Die Fauna des deutschen Unterkarbons, 2 Teil. *Abhandl. Preuss. Geol. Landesanstalt Neve Folge*, **136**, 440 p.
- Pareyn, C., 1961, *Massifs carboniferes du Sahara Sud-Oranais, Tome 1, Publications du Centre de Recherches Sahariennes, serie geologie*, no. 1, Edit. Centre Nation. Recher. Scient., Paris, 202 p.
- Pattison, J., 1981, The stratigraphical distribution of gigantoproductoid brachiopods in the Viséan and Namurian rocks of some areas in northern England. *Rep. Inst. Geol. Sci.*, no. 81/9, 1-30.
- Phillips, J., 1836, *Illustrations of the geology of Yorkshire; or a description of the strata and organic remains: accompanied by a geologic map, sections and diagrams and figures of the fossils, Part 2. The Mountain Limestone Districts*, John Murray, London, 253 p.
- Prentice, J. E., 1956, *Gigantoproductus edelburgensis* (Phillips) and related species. *Proc. Yorkshire geol. Soc.*, **30**, 229-258.
- Qiao, L., and Shen, S.-Z., 2014, Global paleobiogeography of brachiopods during the Mississippian – Response to the global tectonic reconfiguration, ocean circulation, and climate changes. *Gondwana Res.*, **26**, 1173-1185.
- Qiao, L., and Shen, S.-Z., 2015, A global review of the Late Mississippian (Carboniferous) *Gigantoproductus* (Brachiopoda) faunas and the paleogeographical, paleoecological, and paleoclimatic implications. *Palaeogeogr. Palaeoclimat. Palaeoecol.*, **420**, 128-137.
- Sarycheva, T. G., 1928, Lower Moscovian Productids group *Productus giganteus* Mart. (*Gigantella* gen. nov.). *Tr. GNI.*, no. 1, 1-71 (in Russian).
- Sarycheva, T. G., 1960, Order Productida. In Orlov, Y. A., ed., *Basic Paleontologii*, **7**, Akad. Nauk SSSR, Moscow, 221-238 (in Russian).
- Sarycheva, T. G. and Legrand-Blain, M., 1977, Generic composition and evolution of the family Semiplanidae (Brachiopoda). *Paleont. Zhur.*, 1977, no. 2, 70-82 (in Russian).
- Sarycheva, T. G. and Sokolskaya, A. N., 1952, A description of the Paleozoic Brachiopoda of the Moscow Basin. *Tr. Paleont. Inst., Akad. Nauk SSSR*, **38**, 1-307 (in Russian).
- Sarycheva, T. G. and Sokoloskaya, A. N., 1959, On the classification of pseudopunctate brachiopods. *Dok. Akad. Nauk SSSR*, **125**, 181-184 (in Russian).
- Semichatova, S. V., 1975, Brachiopoda. In Garanj, I. M., Guseva, S. N., Devingtal, V. V., Donakova, L. M.,

- Enokyan, N. V., Kalashnikov, N. N., Semichatova, S. V., Stepanov, D. L., Stepanova, G. A., Shostakova, M.F., and Einor, O. L., eds., *Paleontological atlas of Carboniferous strata in the Urals*. Nedra, Leningrad, 154–258 (in Russian).
- Scotese, C. R., 2004, A continental drift flipbook. *J. Geol.*, **112**, 729–741.
- Sowerby, J., 1822, *The Mineral Conchology of Great Britain, vol. 4*, W. Ardling, London, 160 p.
- Stehli, F. G., 1954, Lower Leonardian Brachiopoda of the Sierra Diablo. *Bull. Amer. Mus. Natur. Hist.*, **105**, 257–358.
- Tan, Z., 1987, Brachiopoda. In Regional Geological Survey Party, Hunan Bureau of Geology and Mineral Resources, ed., *Late Devonian and Early Carboniferous stratigraphy and paleontology of Funan*. Geol. Pub. House, Beijing, 111–133 (in Chinese).
- Tazawa, J., 2004, The strike slip model: A synthesis on the origin and tectonic evolution of the Japanese Islands. *Jour. Geol. Soc. Japan*, **110**, 503–517 (in Japanese).
- Tazawa, J., 2018, Early Carboniferous (Mississippian) brachiopods from the Hikoroichi Formation, South Kitakami Belt, Japan. *Mem. Fukui Pref. Dinosaur Mus.*, **17**, 27–87.
- Tazawa, J. and Ibaraki, Y., 2009, *Linoprotonia* and *Gigantoproductus* (Linoproductoidea, Brachiopoda) from the Lower Carboniferous in the Onimaru quarry, Hikoroichi, southern Kitakami Mountains, NE Japan. *Sci. Rep. Niigata Univ., (Geol.)*, no. 24, 7–19.
- Tazawa, J. and Kato, M., 1986, *Striatifera* and *Gigantoproductus* from the Lower Carboniferous of Fukuji, Central Japan. *Trans. Proc. Palaeont. Soc. Japan, N. S.* no. 142, 393–399.
- Tazawa, J., and Miyake, Y., 2002, *Gigantoproductus* (Brachiopoda) from the Lower Carboniferous (Upper Visean) Onimaru Formation of the southern Kitakami Mountains, NE Japan. *Sci. Rep. Niigata Univ., Ser. E*, no. 17, 1–6.
- Tazawa, J., Sato, K. and Takenouchi, K., 2005, *Delepinea* and *Daviesiella* (Chonetoidea, Brachiopoda) from the Lower Carboniferous of Omi, central Japan. *Sci. Rep. Niigata Univ. (Geol.)*, no. 20, 1–13.
- Tazawa, J., and Taira, M., 2020, Early Carboniferous (late Visean) brachiopods from the Onimaru Formation in the Nagaiwa-Onimaru area, South Kitakami Belt, northeastern Japan. *Sci. Rep. Niigata Univ., (Geol.)*, no. 35, 13–31.
- Waagen, W., 1883, Salt Range fossils, 1. Productus-Limestone fossils: Brachiopoda. *Palaeont. Indica, Ser. 13*, **1**, 391–546.
- Wang, S., 1984, Phylum Brachiopoda. In Regional Geological Surveying Team of Hubei, ed., *The Palaeontological Atlas of Hubei Province*. Hubei Sci. Tech. Press, Wuhan, 128–236 (in Chinese).
- Wang, G., Liu, Q., Jing, Y., Hu, S., Liang, W. and Liao, Z., 1982, Brachiopoda. In Nanjin Institute of Geology and Mineral Resources, eds., *Paleontological Atlas of East China (Late Paleozoic)*, **2**, Geol. Pub. House, Beijing, 186–256 (in Chinese).
- Yanagida, J., 1979, The large Carboniferous strophomenides from the Akiyoshi Limestone Group and their biostratigraphic significance. *Proc. Japan Acad., Ser. B*, **55**, 109–114.
- Yanagida, J., 1989, Carboniferous brachiopods from the Akiyoshi Limestone Group, Southwest Japan. In Jin, Y. and Chun, L., eds., *11th Congr. Internat. Strat. Geol. Carbonifere Beijing 1987, Compte Rendu*, **2**, Nanjing Univ. Press, Nanjing, 391–398.
- Yang, Z., Ding, P., Yin, H., Zhabg, S. and Fang, J., 1962, Carboniferous, Permian and Triassic brachiopod faunas from the Chilianshan region. In Chinese Institute of Geology and Palaeontology, Chinese Academy of Sciences, Institute of Geology, Chinese Academy of Sciences, Beijin Institute of Geology, eds., *Monogr. Geol. Chilianshan Mountains*, **4**, Sci. Pub. House, Beijing, 129 p. (in Chinese).
- Yang, S., and Jiang, J., 1987, Early Carboniferous strata and brachiopods of Longmenshan Region, Sichuan. In Bureau of Geology and Mineral Resources of Guizhou Province Commission on Stratigraphy and Palaeontology, Geol. Soc. Guizhou Province, eds., *National Carboniferous Symposium of China, special paper*, Geol. Pub. House, Beijing, 69–92 (in Chinese).
- Yang, D., Ni, S., Chang, M. and Zhao, R., 1977, Brachiopoda. In Hubei Institute of Geological Sciences and others, eds., *Paleontological Atlas of Central-South China, Late Paleozoic Part 2*, Geol. Pub. House, Beijing, 303–470 (in Chinese).
- Zakowa, H., 1986, Brachiopods of the family Semiplanidae Sarycheva, 1960 from the Upper Visean of Poland. *Biul. Inst. Geol.*, no. 355, 49–70.